

BENZOTHAZEPINE DERIVATIVES FOR THE TREATMENT OF HYPERLIPIDEMIA



(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
20 March 2003 (20.03.2003)

PCT WO 03/022825 A1

(10) International Publication Number
WO 03/022825 A1

(51) International Patent Classification: C07D 281/10;
A61K 31/534, C07K 3/06, A61K 38/05, A61P 3/06

(21) International Application Number: PCT/GB02/04043

(74) Agents: BRVANT, Tracey et al.; Astrazeneca, Global Intellectual Property, Mersey, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).

(22) International Filing Date: 5 September 2002 (05.09.2002)

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GR, GU, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Date: 7 September 2001 (07.09.2001) GB 0121621.7

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BI, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TO).


(71) Applicant (for all designated States except MG, US): ASTRAZENECA AB (SE/SE); Sodertalje, S-151 85 Sweden (SE).

(72) Inventors; and
(75) Inventors/Applicants (for US only): STARK, Ingemar (SE/SE); Molndal, S-431 83 Sweden (SE), DAHLSTROM, Mikael, Ulf, Johan (FUSE); Molndal, don W1Y 6LN (GB).

Published:
— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: BENZOTHAZEPINE DERIVATIVES FOR THE TREATMENT OF HYPERLIPIDEMIA



(57) Abstract: The present invention relates to compounds of formula (I), wherein variable groups are as defined within; pharmaceutically acceptable salts, solvates, solvates of such salts and prodrugs thereof and their use as bile acid transport (IBAT) inhibitors for the treatment of hyperlipidemia. Processes for their manufacture and pharmaceutical compositions containing them are also described.

This invention relates to benzothiazepine derivatives, or pharmaceutically acceptable salts, solvates, solvates of such salts and prodrugs thereof. These benzothiazepines possess ileal bile acid transport (IBAT) inhibitory activity and accordingly have value in the treatment of disease states associated with hyperlipidaemic conditions and they are useful in methods of treatment of a warm-blooded animal, such as man. The invention also relates to processes for the manufacture of said benzothiazepine derivatives, to pharmaceutical compositions containing them and to their use in the manufacture of medicaments to inhibit IBAT in a warm-blooded animal, such as man.

It is well-known that hyperlipidaemic conditions associated with elevated concentrations of total cholesterol and low-density lipoprotein cholesterol are major risk factors for cardiovascular atherosclerotic disease (for instance "Coronary Heart Disease: Reducing the Risk; a Worldwide View" Assman G, Cullen P. et al; Circulation 1999, 100, 1930-1938 and "Diabetes and Cardiovascular Disease: A Statement for Healthcare Professionals from the American Heart Association" Grundy S, Benjamin L, Burke G., et al; Circulation, 1999, 100, 1134-46). Interfering with the circulation of bile acids within the lumen of the intestinal tracts is found to reduce the level of cholesterol. Previous established therapies to reduce the concentration of cholesterol involve, for instance, treatment with HMG-CoA reductase inhibitors, preferably statins such as simvastatin and fluvastatin, or treatment with bile acid binders, such as resins. Frequently used bile acid binders are for instance colestyramine and colestipol. One recently proposed therapy ("Bile Acids and Lipoprotein Metabolism: a Renaissance for Bile Acids in the Post Statin Era" Angelin B, Eriksson M, Rudling M; Current Opinion on Lipidology, 1999, 10, 269-74) involved the treatment with substances with an IBAT inhibitory effect.


Re-absorption of bile acid from the gastro-intestinal tract is a normal physiological process which mainly takes place in the ileum by the IBAT mechanism. Inhibitors of IBAT can be used in the treatment of hypercholesterolaemia (see for instance "Interaction of bile acids and cholesterol with nonsystemic agents having hypocholesterolaemic properties", Biochemica et Biophysica Acta, 1210 (1994) 255-287). Thus, suitable compounds having such inhibitory IBAT activity are also useful in the treatment of hyperlipidaemic conditions. Compounds possessing such IBAT inhibitory activity have been described, see for instance the compounds described in WO 93/16055, WO 94/18183, WO 94/18184, WO 96/05188,

WO 96/08488, WO 96/16051, WO 97/33882, WO 98/38182, WO 99/35153, WO 98/40375, WO 99/35153, WO 99/64409, WO 99/64410, WO 00/01687, WO 00/47568, WO 00/61568, WO 01/68906, DE 19825804, WO 00/38725, WO 00/38726, WO 00/38727, WO 00/38728, WO 00/38729, WO 01/68906, WO 01/66553, WO 02/50051 and EP 0 864 582.

A further aspect of this invention relates to the use of the compounds of the invention in the treatment of dyslipidemic conditions and disorders such as hypertriglyceridemia, hypertriglyceridemia, hyperbetaipoproteinemia (high LDL), hypertriglyceridemia (high VLDL), hyperchylomicronemia, hypolipoproteinemia, hypercholesterolemia, hyperlipoproteinemia and hypoalphalipoproteinemia (low HDL). In addition, these compounds are expected to be useful for the prevention and treatment of different clinical conditions such as atherosclerosis, arteriosclerosis, arrhythmia, hyper-thrombotic conditions, vascular dysfunction, endothelial dysfunction, heart failure, coronary heart diseases, cardiovascular diseases, myocardial infarction, angina pectoris, peripheral vascular diseases, inflammation of cardiovascular tissues such as heart, valves, vasculature, arteries and veins, aneurisms, stenosis, restenosis, vascular plaques, vascular fatty streaks, leukocytes, monocytosis and/or macrophage infiltration, intimal thickening, medial thinning, infectious and surgical trauma and vascular thrombosis, stroke and transient ischemic attacks.

The present invention is based on the discovery that certain benzothiazepine compounds surprisingly inhibit IBAT. Such properties are expected to be of value in the treatment of disease states associated with hyperlipidemic conditions.

Accordingly, the present invention provides a compound of formula (I):




wherein:

25 One of R^1 and R^2 are selected from hydrogen, C_1 -alkyl or C_2 -alkenyl and the other is selected from C_1 -alkyl or C_2 -alkenyl;

R¹ is selected from hydrogen, hydroxy, C₁-alkyl, C₁-alkoxy and C₁-alkenylloxy;
R² is selected from halo, nitro, cyano, hydroxy, amino, carbony, carbamoyl, mercapto, sulphanoyl, C₁-alkyl, C₂-alkenyl, C₂-alkynyl, C₁-alkoxy, C₁-alkenoyl, C₁-alkenylloxy, *N*-(C₁-alkyl)amino, *N,N'*-(C₁-alkyl)₂amino, C₁-alkenylamino, *N'*-(C₁-alkyl)carbamoyl, *N,N'*-(C₁-alkyl)₂carbamoyl, C₁-alkylSi(O)_n wherein *n* is 0 to 2, C₁-alkoxycarbonyl, *N*-(C₁-alkyl)sulphanoyl and *N,N'*-(C₁-alkyl)₂sulphanoyl;

one of R^4 and R^5 is a group of formula (IA):



10


R² and **R⁶** and the other of **R⁴** and **R⁵** are independently selected from hydrogen, halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphanoyl, C₁-alkyl, C₂-alkenyl, C₂-alkynyl, C₁-alkoxy, C₁-alkenyl, C₁-alkanyloxy, N-(C₁-alkyl)amino, N,N-(C₁-alkyl)₂amino, C₁-alkanoylamino, N-(C₁-alkyl)carbamoyl, N,N-(C₁-alkyl)₂carbamoyl, C₁-alkylS(O)_n wherein n is 0 to 2, C₁-alkoxycarbonyl, N-(C₁-alkyl)sulphanoyl and N,N-(C₁-alkyl)₂sulphanoyl; wherein R³ and R⁵ and the other of R⁴ and R⁵ may be optionally substituted on carbon by one or more R¹⁶,

2;
20 Ring A is aryl or heteroaryl, wherein Ring A is optionally substituted by one or more substituents selected from R¹⁷;

R¹ is hydrogen, C₁-alkyl, carbocycyl or heterocycyl; wherein R is optionally substituted by one or more substituents selected from R¹⁸;

R⁸ is hydrogen or C₁₋₄alkyl;
R⁹ is hydrogen or C₁₋₄alkyl;
R¹⁰ is hydrogen, C₁₋₄alkyl, carbocyclyl or heterocyclyl, wherein R¹⁰ is optionally substituted by one or more substituents selected from R¹⁹;

R^{11} is carboxy, sulpho, sulphino, phosphono, $-P(O)(OR^b)(OR^b)$, $-P(O)(OH)(OR^b)$, $-P(O)(OH)(R^b)$ or $-P(O)(OR^b)(R^b)$ wherein R^b and R^d are independently selected from C_1 -alkyl; or R^{11} is a group of formula (IB):




wherein:

Y is $-N(R^3)$, $-N(R^3)C(O)$, $-O-$, and $-S(O)_a$; wherein a is 0-2 and R^2 is hydrogen or C_1 -alkyl;

R^{12} is hydrogen or C_1 -alkyl;

R^{13} and R^{14} are independently selected from hydrogen, C_1 -alkyl, carbocyclyl or heterocyclyl; wherein R^{13} and R^{14} may be independently optionally substituted by one or more substituents selected from R^{20} ;

R^{15} is carboxy, sulpho, sulphino, phosphono, $-P(O)(OR^b)(OR^b)$, $-P(O)(OH)(OR^b)$, $-P(O)(OH)(R^b)$ or $-P(O)(OR^b)(R^b)$ wherein R^b and R^f are independently selected from C_1 -alkyl; or R^{15} is a group of formula (IC):



wherein:

R^{24} is selected from hydrogen or C_1 -alkyl;

R^{25} is selected from hydrogen, C_1 -alkyl, carbocyclyl, heterocyclyl or R^{27} ; wherein said C_1 -alkyl, carbocyclyl or heterocyclyl may be independently optionally substituted by one or more substituents selected from R^{28} ;

R^{26} is selected from carboxy, sulpho, sulphino, phosphono, tetrazolyl,

$-P(O)(OR^b)(OR^b)$, $-P(O)(OH)(OR^b)$, $-P(O)(OH)(R^b)$ or $-P(O)(OR^b)(R^b)$ wherein R^b and R^g are independently selected from C_1 -alkyl;

p is 1-3; wherein the values of R^{13} may be the same or different;

q is 0-1;

r is 0-3; wherein the values of R^{14} may be the same or different;
 m is 0-2; wherein the values of R^{10} may be the same or different;
 n is 1-3; wherein the values of R^7 may be the same or different;

z is 0-3; wherein the values of R^{23} may be the same or different;


R^{16} , R^{17} and R^{18} are independently selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphonoyl, C_1 -alkyl, C_2 -alkenyl, C_2 -alkynyl, C_1 -alkoxy, C_1 -alkanoyl, C_1 -alkanoyloxy, N -(C_1 -alkyl)amino, N,N -(C_1 -alkyl)₂amino, C_1 -alkanoylamino, N -(C_1 -alkyl)carbamoyl, N,N -(C_1 -alkyl)₂carbamoyl, C_1 -alkylS(O)₂, wherein a is 0 to 2, C_1 -alkoxy, carbonyl, N -(C_1 -alkyl)sulphonoyl and N,N -(C_1 -alkyl)₂sulphonoyl; wherein R^{16} , R^{17} and R^{18} may be independently optionally substituted on carbon by one or more R^{21} ;

R^{19} , R^{20} , R^{27} and R^{28} are independently selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphonoyl, C_1 -alkyl, C_2 -alkenyl, C_2 -alkynyl, C_1 -alkoxy, C_1 -alkanoyl, C_1 -alkanoyloxy, N -(C_1 -alkyl)amino, N,N -(C_1 -alkyl)₂amino, C_1 -alkanoylamino, N -(C_1 -alkyl)carbamoyl, N,N -(C_1 -alkyl)₂carbamoyl, C_1 -alkylS(O)₂, wherein a is 0 to 2, C_1 -alkoxy, carbonyl, N -(C_1 -alkyl)sulphonoyl, N,N -(C_1 -alkyl)₂sulphonoyl, carbocyclyl, heterocyclyl, sulpho, sulphino, amidino, (C_1 -alkyl)silyl, phosphono, $-P(O)(OR^b)(OR^b)$, $-P(O)(OH)(OR^b)$, $-P(O)(OH)(R^b)$ or $-P(O)(OR^b)(R^b)$, wherein R^b and R^b are independently selected from C_1 -alkyl; wherein R^{19} and R^{20} may be independently optionally substituted on carbon by one or more R^{22} ;

R^{21} and R^{22} are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carboxy, carbamoyl, mercapto, sulphonoyl, trifluoromethyl, trifluoromethoxy, methyl, ethyl, methoxy, ethoxy, vinyl, allyl, ethynyl, methoxycarbonyl, formyl, acetyl, formamido, acetylamino, acetoxycarbonyl, methoxy, methylamino, dimethylamino, N -methylcarbamoyl, N,N -dimethylcarbamoyl, methylthio, methylsulphonyl, mesyl, N -methylsulphonoyl and N,N -dimethylsulphonoyl;

According to a further feature of the present invention there is provided a compound of formula (I):

- 6 -



wherein:

5 One of R^1 and R^2 are selected from hydrogen or C_{1-6} alkyl and the other is selected from C_{1-6} alkyl;

R^2 is selected from hydrogen, hydroxy, C_{1-6} alkyl, C_{1-6} alkoxy and C_{1-6} alkanoyloxy;

R^3 is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbonyl, mercapto,

sulphamoyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy,


$N-(C_{1-6}alkyl)amino$, $N,N-(C_{1-6}alkyl)_2amino$, $C_{1-6}alkanoylamino$, $N-(C_{1-6}alkyl)carbamoyl$,

10 $N,N-(C_{1-6}alkyl)_2carbamoyl$, $C_{1-6}alkylS(O)_2$, wherein a is 0 to 2, $C_{1-6}alkoxy$ carbonyl,

$N-(C_{1-6}alkyl)sulphamoyl$ and $N,N-(C_{1-6}alkyl)_2sulphamoyl$;

v is 0-5;

one of R^4 and R^5 is a group of formula (IA):



15 (IA)

R^3 and R^6 and the other of R^4 and R^5 are independently selected from hydrogen, halo,

nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, C_{1-6} alkyl,

C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, $N-(C_{1-6}alkyl)amino$,

$N,N-(C_{1-6}alkyl)_2amino$, $C_{1-6}alkanoylamino$, $N-(C_{1-6}alkyl)carbamoyl$,

20 $N,N-(C_{1-6}alkyl)_2carbamoyl$, $C_{1-6}alkylS(O)_2$, wherein a is 0 to 2, $C_{1-6}alkoxy$ carbonyl,

$N-(C_{1-6}alkyl)sulphamoyl$ and $N,N-(C_{1-6}alkyl)_2sulphamoyl$; wherein R^3 and R^6 and the other of

R^4 and R^5 may be optionally substituted on carbon by one or more R^{16} ;

- 7 -

2; X is O , $N(R^7)$, $S(O)_b$ or $-CH(R^7)$; wherein R^7 is hydrogen or C_{1-6} alkyl and b is 0-

Ring A is aryl or heteroaryl; wherein Ring A is optionally substituted by one or more

substituents selected from R^{17} ;

5 R^7 is hydrogen, C_{1-6} alkyl, carbocyclyl or heterocyclyl; wherein R^7 is optionally

substituted by one or more substituents selected from R^{18} ;

R^8 is hydrogen or C_{1-6} alkyl;

R^9 is hydrogen or C_{1-6} alkyl;


R^{10} is hydrogen, C_{1-6} alkyl, carbocyclyl or heterocyclyl; wherein R^{10} is optionally

0 substituted by one or more substituents selected from R^{19} ;

R^{11} is carboxy, sulphy, sulphino, phosphono, $-P(O)(OR^9)(OR^9)$, $-P(O)(OH)(OR^9)$,

$-P(O)(OH)(R^9)$ or $-P(O)(OR^9)(R^9)$ wherein R^9 and R^9 are independently selected from

C_{1-6} alkyl; or R^{11} is a group of formula (IB):



15 (IB)

wherein:

Y is $-N(R^9)$, $-N(R^9)C(O)_2$, $-O_2$ and $-S(O)_a$; wherein a is 0-2 and R^9 is hydrogen or

C_{1-6} alkyl;

R^{12} is hydrogen or C_{1-6} alkyl;

20 R^{13} and R^{14} are independently selected from hydrogen, C_{1-6} alkyl, carbocyclyl or

heterocyclyl; wherein R^{13} and R^{14} may be independently optionally substituted by one or more

substituents selected from R^{20} ;

R^{15} is carboxy, sulphy, sulphino, phosphono, $-P(O)(OR^9)(OR^9)$, $-P(O)(OH)(OR^9)$,

$-P(O)(OH)(R^9)$ or $-P(O)(OR^9)(R^9)$ wherein R^9 and R^9 are independently selected from C_{1-6} alkyl

25 p is 1-3; wherein the values of R^{13} may be the same or different;

q is 0-1;

r is 0-3; wherein the values of R^{14} may be the same or different;

m is 0-2; wherein the values of R^{10} may be the same or different;

n is 1-3; wherein the values of R^7 may be the same or different;

R^{16} , R^{17} and R^{18} are independently selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphanoyl, C_1 -alkyl, C_2 -alkenyl, C_2 -alkynyl, C_1 -alkoxy, C_1 -alkanoyl, C_1 -alkanoyloxy, N -(C_1 -alkyl)amino, N,N -(C_1 -alkyl) $_2$ amino, C_1 -alkanoylamino, N -(C_1 -alkyl)carbamoyl, N,N -(C_1 -alkyl) $_2$ carbamoyl, C_1 -alkylS(O), wherein a is 0 to 2, C_1 -alkoxy-carbonyl, M -(C_1 -alkyl)sulphanoyl and N,N -(C_1 -alkyl) $_2$ sulphanoyl; wherein R^{16} , R^{17} and R^{18} may be independently optionally substituted on carbon by one or more R^{21} .

R^{19} and R^{20} are independently selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphanoyl, C_1 -alkyl, C_2 -alkenyl, C_2 -alkynyl, C_1 -alkoxy, C_1 -alkanoyl, C_1 -alkanoyloxy, N -(C_1 -alkyl)amino, N,N -(C_1 -alkyl) $_2$ amino, C_1 -alkanoylamino, N -(C_1 -alkyl)carbamoyl, N,N -(C_1 -alkyl) $_2$ carbamoyl, C_1 -alkylS(O), wherein a is 0 to 2, C_1 -alkoxy-carbonyl, M -(C_1 -alkyl)sulphanoyl, N,N -(C_1 -alkyl) $_2$ sulphanoyl, carbocyclyl, heterocyclyl, sulpho, sulphino, amidino, phosphono, $-P(O)(OR^a)(OR^b)$, $-P(O)(OH)(OR^a)$, $-P(O)(OH)(R^a)$ or $-P(O)(OR^a)(R^b)$, wherein R^a and R^b are independently selected from C_1 -alkyl; wherein R^{19} and R^{20} may be independently optionally substituted on carbon by one or more R^{22} .

R^{21} and R^{22} are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carboxy, carbamoyl, mercapto, sulphanoyl, trifluoromethyl, trifluoromethoxy, methyl, ethyl, methoxy, ethoxy, vinyl, allyl, ethynyl, methoxy-carbonyl, formyl, acetyl, formamido, acetyl/amino, acetoxymethyl, methyl/amino, dimethyl/amino, N -methylcarbamoyl, N,N -dimethylcarbamoyl, methylthio, methylsulphinyl, mesyl, N -methylsulphanoyl and N,N -dimethylsulphanoyl;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

In this specification the term "alkyl" includes both straight and branched chain alkyl groups but references to individual alkyl groups such as "propyl" are specific for the straight chain version only. For example, " C_1 -alkyl" includes C_1 -alkyl, propyl, isopropyl and t -butyl. However, references to individual alkyl groups such as 'propyl' are specific for the straight chained version only and references to individual branched chain alkyl groups such as 'isopropyl' are specific for the branched chain version only. A similar convention applies to other radicals, for example "phenyl/ C_1 -alkyl" would include phenyl/ C_1 -alkyl, benzyl, 1-phenylethyl and 2-phenylethyl. The term "halo" refers to fluoro, chloro, bromo and iodo.

Where optional substituents are chosen from "one or more" groups it is to be understood that this definition includes all substituents being chosen from one of the specified groups or the substituents being chosen from two or more of the specified groups.

"Heteroaryl" is a totally unsaturated, mono or bicyclic ring containing 3-12 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked. Preferably "heteroaryl" refers to a totally unsaturated, monocyclic ring containing 5 or 6 atoms or a bicyclic ring containing 9 or 10 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked. Examples and suitable values of the term "heteroaryl" are thienyl, isoxazolyl, imidazolyl, pyrrolyl, thiadiazolyl, isothiazolyl, triazolyl, pyranyl, indolyl, pyrimidyl, pyrazinyl, pyridazinyl, pyridyl and quinolyl. Preferably the term "heteroaryl" refers to thienyl or indolyl.

"Aryl" is a totally unsaturated, mono or bicyclic carbon ring that contains 3-12 atoms. Preferably "aryl" is a monocyclic ring containing 5 or 6 atoms or a bicyclic ring containing 9 or 10 atoms. Suitable values for "aryl" include phenyl or naphthyl. Particularly "aryl" is phenyl.

A "heterocyclyl" is a saturated, partially saturated or unsaturated, mono or bicyclic ring containing 3-12 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked, wherein a $-CH_2-$ group can optionally be replaced by a $-C(O)-$ or a ring sulphur atom may be optionally oxidised to form the S-oxides. Preferably a "heterocyclyl" is a saturated, partially saturated or unsaturated, mono or bicyclic ring containing 5 or 6 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked, wherein a $-CH_2-$ group can optionally be replaced by a $-C(O)-$ or a ring sulphur atom may be optionally oxidised to form S-oxide(s). Examples and suitable values of the term "heterocyclyl" are thiazolidinyl, pyrrolidinyl, pyrrolinyl, 2-pyrrolidinyl, 2,5-dioxypyrrolidinyl, 2-benzoxazolinonyl, 1,1-dioxotetrahydrothienyl,

2,4-dioxoimidazolidinyl, 2-oxo-1,3,4-(4-triazolinyl), 2-oxazolidinonyl, 5,6-dihydrouracilyl, 1,3-benzodioxolyl, 1,2,4-oxadiazolyl, 2-azabicyclo[2.2.1]heptyl, 4-thiazolidinonyl, morpholino, 2-oxotetrahydrofuranyl, tetrahydrofuranyl, 2,3-dihydrobenzofuranyl, benzothienyl, tetrahydropyranyl, piperidyl, 1-oxo-1,3-dihydroisindolyl, piperazinyl, thiomorpholino, 1,1-dioxothiomoorpholino, tetrahydropyranyl, 1,3-dioxolanyl, homopiperazinyl, thienyl, isoxazolyl, imidazolyl, pyrrolyl, thiadiazolyl, isothiazolyl, 1,2,4-triazolyl, 1,3,4-triazolyl,

pyranyl, indolyl, pyrimidyl, thiazolyl, pyrazinyl, pyridazinyl, pyridyl, 4-pyridonyl, quinolyl and 1-isouquinolonyl.

A "carbocyclyl" is a saturated, partially saturated or unsaturated, mono or bicyclic carbon ring that contains 3-12 atoms, wherein a -CH₂- group can optionally be replaced by a -C(O)-. Preferably "carbocyclyl" is a monocyclic ring containing 5 or 6 atoms or a bicyclic ring containing 9 or 10 atoms. Suitable values for "carbocyclyl" include cyclopropyl, cyclobutyl, 1-oxocyclopentyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, phenyl, naphthyl, tetralinyl, indenyl or 1-oxoindanyl. Particularly "carbocyclyl" is cyclopropyl, cyclobutyl, 1-oxocyclopentyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, phenyl or 1-oxoindanyl.

An example of "C₁-alkamoyloxy" and "C₁-alkamoyloxy" is acetoxy. Examples of "C₁-alkoxycarbonyl" and "C₁-alkoxycarbonyl" include methoxycarbonyl, ethoxycarbonyl, *n*- and *i*-butoxycarbonyl. Examples of "C₁-alkoxy" "C₁-alkoxy" include methoxy, ethoxy and propoxy. Examples of "C₁-alkanoylamino" and "C₁-alkanoylamino" include formamido, acetamido and propionylamino. Examples of "C₁-alkylS(O)_n" wherein *n* is 0 to 2" and "C₁-alkylS(O)_n" wherein *n* is 0 to 2" include methylthio, ethylthio, methylsulphonyl, ethylsulphonyl, mesyl and ethylsulphonyl. Examples of "C₁-alkanoyl" and "C₁-alkanoyl" include propionyl and acetyl. Examples of "N-(C₁-alkyl)amino" and "N-(C₁-alkyl)amino" include methylamino and ethylamino. Examples of "N,N-(C₁-alkyl)₂amino" and "N,N-(C₁-alkyl)₂amino" include di-*N*-methylamino, di-*N*-ethylamino and *N*-ethyl-*N*-methylamino. Examples of "C₂-alkenyl" and "C₂-alkenyl" are vinyl, allyl and 1-propenyl. Examples of "C₂-alkynyl" and "C₂-alkynyl" are ethynyl, 1-propynyl and 2-propynyl. Examples of "N-(C₁-alkyl)sulphamoyl" and "N-(C₁-alkyl)sulphamoyl" are *N*-(methyl)sulphamoyl and *N*-(ethyl)sulphamoyl. Examples of "N-(C₁-alkyl)sulphamoyl" and "N-(C₁-alkyl)sulphamoyl" are *N,N*-(dimethyl)sulphamoyl and *N,N*-(dimethyl)sulphamoyl and *N*-(methyl)-*N*-(ethyl)sulphamoyl. Examples of "N-(C₁-alkyl)carbamoyl" and "N-(C₁-alkyl)carbamoyl" are methylaminocarbonyl and ethylaminocarbonyl. Examples of "N,N-(C₁-alkyl)₂carbamoyl" and "N,N-(C₁-alkyl)₂carbamoyl" are dimethylaminocarbonyl and methyl ethylaminocarbonyl. Examples of "(C₁-alkyl)silyl" include trimethylsilyl and methyl diethylsilyl.

A suitable pharmaceutically acceptable salt of a compound of the invention is, for example, an acid-addition salt of a compound of the invention which is sufficiently basic, for example, an acid-addition salt with, for example, an inorganic or organic acid, for example

hydrochloric, hydrotromic, sulphuric, phosphoric, trifluoroacetic, citric, acetate or maleic acid. In addition a suitable pharmaceutically acceptable salt of a compound of the invention which is sufficiently acidic is an alkali metal salt, for example a sodium or potassium salt, an alkaline earth metal salt, for example a calcium or magnesium salt, an ammonium salt or a salt with an organic base which affords a physiologically-acceptable cation, for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or *tert*-(2-hydroxyethyl)amine.

The compounds of the formula (I) may be administered in the form of a pro-drug which is broken down in the human or animal body to give a compound of the formula (I), examples of pro-drugs include *in vivo* hydrolysable esters and *in vivo* hydrolysable amides of a compound of the formula (I).

An *in vivo* hydrolysable ester of a compound of the formula (I) containing carboxy or hydroxy group is, for example, a pharmaceutically acceptable ester which is hydrolysed in the human or animal body to produce the parent acid or alcohol. Suitable pharmaceutically acceptable esters for carboxy include C₁-alkoxymethyl esters for example methoxymethyl, C₁-alkanoyloxymethyl esters for example pivaloyloxymethyl, phthalidyl esters, C₃-acyloalkoxycarbonyloxycarbonyl esters for example 1-cyclohexylcarbonyloxycarbonyl, 1,3-dioxolan-2-onylmethyl esters for example 5-methyl-1,3-dioxolan-2-onylmethyl, and C₁-alkoxycarbonyloxycarbonyl esters for example 1-methoxycarbonyloxycarbonyl and may be formed at any carboxy group in the compounds of this invention.

An *in vivo* hydrolysable ester of a compound of the formula (I) containing a hydroxy group includes inorganic esters such as phosphate esters and α -acyloxyalkyl ethers and related compounds which as a result of the *in vivo* hydrolysis of the ester breakdown to give the parent hydroxy group. Examples of α -acyloxyalkyl ethers include acetoxy methoxy and 2,2-dimethylpropionyloxy-methoxy. A selection of *in vivo* hydrolysable ester forming groups for hydroxy include alkamoyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl, alkoxycarbonyl (to give alkyl carbonate esters), dialkylcarbamoyl and *N*-(dialkylaminomethyl)-*N*-alkylcarbamoyl (to give carbamates), dialkylaminocarbonyl and carboxyacetate. Examples of substituents on benzoyl include morpholino and piperazino linked from a ring nitrogen atom via a methylene group to the 3- or 4- position of the benzoyl ring.

A suitable value for an *in vivo* hydrolysable amide of a compound of the formula (I) containing a carboxy group is, for example, a *N*-C₁₋₆alkyl or *N,N*-di-C₁₋₆alkyl amide such as *N*-methyl, *N*-ethyl, *N*-propyl, *N,N*-dimethyl, *N*-ethyl-*N*-methyl or *N,N*-diethyl amide.

Some compounds of the formula (I) may have chiral centres and/or geometric isomeric centres (*E*- and *Z*- isomers), and it is to be understood that the invention encompasses all such optical, diastereoisomers and geometric isomers that possess IBAT inhibitory activity.

The invention relates to any and all tautomeric forms of the compounds of the formula (I) that possess IBAT inhibitory activity.

It is also to be understood that certain compounds of the formula (I) can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms which possess IBAT inhibitory activity.

Particular values are as follows. Such values may be used where appropriate with any of the definitions, claims or embodiments defined hereinbefore or hereinafter.

15 R¹ and R² are C₁-alkyl.

R¹ and R² are butyl.

One of R¹ and R² is ethyl and the other is butyl.

One of R¹ and R² is ethyl and the other is butyl or R¹ and R² are both butyl.

v is 0 or 1.

v is 0.

R² is C₁-alkyl.

R² is hydrogen.

R² is hydrogen or hydroxy.

R³ and R⁶ are hydrogen.

R⁴ is methylthio.

R⁴ is hydrogen.

R⁴ is hydrogen, halo or C₁-alkyl(S(O)_n), wherein n is 0.

R⁴ is hydrogen, bromo or methylthio.

R² is a group of formula (IA) (as depicted above) wherein:

X is -O-;

n is 1;

R² is hydrogen;

R⁸ is hydrogen;

R⁹ is hydrogen;

m is 0; and

R¹¹ is carboxy.

5 R² is *N*-(R)- α -carboxybenzyl)carbamoylmethoxy.

R² is a group of formula (IA) (as depicted above); wherein

X is -O-;

Ring A is aryl; wherein Ring A is optionally substituted by one or more substituents selected from R¹⁷;

10 R⁷ is hydrogen;

R⁸ is hydrogen;

R⁹ is hydrogen;

R¹¹ is carboxy; or R¹¹ is a group of formula (IB) (as depicted above); wherein:

R¹² is hydrogen;

15 R¹³ is hydrogen;

R¹⁵ is carboxy or sulpho;

p is 1 or 2;

q is 0;

r is 0;

20 m is 0;

n is 1; and

R¹⁷ is hydroxy.

R² is a group of formula (IA) (as depicted above); wherein

X is -O-;

25 Ring A is phenyl or 4-hydroxyphenyl;

R⁷ is hydrogen;

R⁸ is hydrogen;

R⁹ is hydrogen;

R¹¹ is carboxy; or R¹¹ is a group of formula (IB) (as depicted above); wherein:

30 R¹² is hydrogen;

R¹³ is hydrogen;

R¹⁵ is carboxy or sulpho;

p is 1 or 2;

q is O₂;

r is O₂;

m is O₂, and

n is 1;

R⁵ is N-((R)-(R)-α-carboxybenzyl)carbamoylmethoxy; N-((R)-(R)-α-[N-(carboxymethyl)carbamoyl]benzyl)carbamoylmethoxy; or N-((R)-(R)-α-[N-(2-sulphoethyl)carbamoyl]-4-hydroxybenzyl)carbamoylmethoxy.

Therefore in one aspect of the invention there is provided a compound of formula (I)

(as depicted above)

R¹ and R² are C₁₋₆alkyl;

v is O₂;

R⁷ is hydrogen or hydroxy;

R³ and R⁶ are hydrogen;

R⁴ is hydrogen, halo or C₁₋₆alkylS(O)_n, wherein n is 0;

R⁵ is a group of formula (IA) (as depicted above); wherein

X is -O-;

Ring A is aryl; wherein Ring A is optionally substituted by one or more substituents selected from R¹⁷;

R⁷ is hydrogen;

R⁸ is hydrogen;

R⁹ is hydrogen;

R¹¹ is carboxy; or R¹¹ is a group of formula (IB) (as depicted above); wherein:

R¹² is hydrogen;

R¹³ is hydrogen;

R¹⁵ is carboxy or sulpho;

p is 1 or 2;

q is O₂;

r is O₂;

m is O₂;

n is 1; and

R¹⁷ is hydroxy;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

Therefore in one aspect of the invention there is provided a compound of formula (I)

(as depicted above)

One of R¹ and R² is ethyl and the other is butyl;

v is O₂;

R⁷ is hydrogen or hydroxy;

R³ and R⁶ are hydrogen;

R⁴ is hydrogen, bromo or methylthio;

R⁵ is N-((R)-(R)-α-carboxybenzyl)carbamoylmethoxy; N-((R)-(R)-α-[N-(carboxymethyl)carbamoyl]benzyl)carbamoylmethoxy; or N-((R)-(R)-α-[N-(2-sulphoethyl)carbamoyl]-4-hydroxybenzyl)carbamoylmethoxy;

(carboxymethyl)carbamoyl]benzyl)carbamoylmethoxy; or N-((R)-(R)-α-[N-(2-sulphoethyl)carbamoyl]-4-hydroxybenzyl)carbamoylmethoxy;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.


In another aspect of the invention, preferred compounds of the invention are any one of the examples or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

15 prodrug thereof.

Preferred aspects of the invention are those which relate to the compound of formula (I) or a pharmaceutically acceptable salt thereof.


Another aspect of the present invention provides a process for preparing a compound of formula (I) or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof which process (wherein variable groups are, unless otherwise specified, as defined in formula (I)) comprises of:

Process 1): oxidising a benzothiazepine of formula (II):




(II);

25 Process 2): for compounds of formula (I) wherein X is -O-, -NR⁸ or -S-, reacting a compound of formula (IIIa) or (IIIb):




with a compound of formula (IV):




wherein L is a displaceable group;


Process 3): reacting an acid of formula (Va) or (Vb):




10 or an activated derivative thereof, with an amine of formula (VI):




Process 4): for compounds of formula (I) wherein R¹¹ is a group of formula (IB); reacting a compound of formula (I) wherein R¹¹ is carboxy with an amine of formula (VII):



5 Process 5): for compounds of formula (I) wherein R¹¹ is a group of formula (IB) and R¹⁵ is a group of formula (IC) reacting a compound of formula (I) wherein R¹⁵ is carboxy with an amine of formula (VIII):



10 Process 6) for compounds of formula (I) wherein one of R⁶ and R⁷ are independently selected from C₁₋₆alkylthio optionally substituted on carbon by one or more R¹⁶, reacting a compound of formula (IXa) or (IXb):




15 wherein L is a displaceable group; with a thiol of formula (X):




wherein R^m is C₁₋₆alkylthio optionally substituted on carbon by one or more R¹⁶;

Process 7): for compounds of formula (I) wherein R¹¹ is carboxy; deprotecting a compound of formula (XIa):



(XIa)


or (XIb):



(XIb)

wherein R^2 together with the $-OC(O)-$ group to which it is attached forms an ester,
Process 8: for compounds of formula (I) wherein R^{11} is a group of formula (IB) and R^{15} is


carboxy, denoting a compound of formula (XIIa):



(XIIa)


or (XIIb):

10




(XIIIa)

wherein R^2 together with the $-OC(O)-$ group to which it is attached forms an ester,
Process 9: for compounds of formula (I) wherein R^{11} is a group of formula (IB) and Y is
 $-N(R^3)C(O)-$ reacting an acid of formula (XIIIb):




(XIIIa)

or (XIIIb):




(XIIIb)

or an activated derivative thereof, with an amine of formula (XIV):




or *Process 10*): for compounds of formula (I) wherein R^{11} is a group of formula (IB), R^{15} is a group of formula (IC) and R^{26} is carboxy; deprotecting a compound of formula (XVa):



(XVa)

or (XVb):



(XVb)

wherein R^1 together with the $-OC(O)-$ group to which it is attached forms an ester, and thereafter if necessary or desirable:

- converting a compound of the formula (I) into another compound of the formula (I);
- removing any protecting groups;
- forming a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug.

L is a displaceable group, suitable values for L are for example, a halogeno or sulphonyloxy group, for example a chloro, bromo, methanesulphonyloxy or


toluene-4-sulphonyloxy group.

R^1 together with the $-OC(O)-$ group to which it is attached forms an ester. Preferably R^1 is methyl or ethyl. More preferably R^1 is methyl. In another aspect of the invention R^1 is C_{1-6} alkyl or phenyl/ C_{1-6} alkyl, preferably C_{1-6} alkyl or benzyl, more preferably *t*-butyl, methyl, ethyl or benzyl.

Specific reaction conditions for the above reactions are as follows.

Process 1): Benzothiazepines of formula (II) may be oxidised under standard sulphur oxidation conditions, for example using hydrogen peroxide and trifluoroacetic acid at a temperature in the range of 0°C to reflux, preferably at or near room temperature.

Compounds of formula (II) may be prepared according to Scheme I:



Scheme 1

Compounds of formula (IIa), (IIb) and (IIc) are commercially available compounds, or they are known in the literature, or they are prepared by standard processes known in the art.

Process 2: Compounds of formula (IIa) or (IIb) may be reacted with compounds of formula (IV) in the presence of a base for example an inorganic base such as sodium carbonate, or an organic base such as Hunigs base, in the presence of a suitable solvent such as acetonitrile, dichloromethane or tetrahydrofuran at a temperature in the range of 0°C to reflux, preferably at or near reflux.


Compounds of formula (IIa) or (IIb) may be prepared in a similar manner to compounds of formula (II) but wherein R⁴ or R⁵ is -OH, -NH(R⁹) or -SH (optionally for -SO₂ followed by the oxidation step of Process 1). Compounds of formula (IIa) or (IIb) wherein X is -O- or -S- may also be prepared by the procedures disclosed in WO 9605188.

Compounds of formula (IV) are commercially available compounds, or they are known in the literature, or they are prepared by standard processes known in the art.

Process 3) and Process 4) and Process 5): Acids and amines may be coupled together in the presence of a suitable coupling reagent. Standard peptide coupling reagents known in the art can be employed as suitable coupling reagents, or for example carbonyldiimidazole and diisobutylcarbodiimide, optionally in the presence of a catalyst such as dimethylaminopyridine or 4-pyrrolidinopyridine, optionally in the presence of a base for example triethylamine, pyridine, or 2,6-di-*tert*-butyl-pyridines such as 2,6-lutidine or benzene, tetrahydrofuran and dimethylformamide. The coupling reaction may conveniently be performed at a temperature in the range of -40 to 40°C.

Suitable activated acid derivatives include acid halides, for example acid chlorides, and active esters, for example pentafluorophenyl esters. The reaction of these types of compounds with amines is well known in the art, for example they may be reacted in the presence of a base, such as those described above, and in a suitable solvent, such as those described above. The reaction may conveniently be performed at a temperature in the range of -40 to 40°C.


Compounds of formula (Va) or (Vb) wherein X=O, -NR⁺, -S- may be prepared according to Scheme 2:



Scheme 2

Wherein L in (IXa) and (IXb) is a displaceable group e.g. bromo, chloro, fluoro, mesyl or tosyl and wherein X is -O-, -S-, NR' (optionally for -SO- and -SO₂- followed by the oxidation step of Process 1).

Compounds of formula (Va) and (Vb) where X is -SO- or -SO₂- may be prepared by oxidising the resulting compounds of formula (Va) and (Vb) from Scheme 2 where X is -S-. Compounds of formula (Va) or (Vb) wherein X is -CH₂-, and n is 1, may be prepared according to Scheme 3.



Scheme 3

The skilled person will appreciate that the above reaction scheme may be manipulated to prepare compounds of formula (Va) or (Vb) where n is 2 or 3.

Compounds of formula (XIIa) or (XIIb) may be prepared by manipulations known to the skilled person of the processes described herein.

Compounds of formula (Ve), (Vf), (VII), (VIII) and (XIV) are commercially available compounds, or they are known in the literature, or they are prepared by standard processes known in the art.

Process 6: Compounds of formula (IXa) and (IXb) may be reacted with thiols of formula (X) in the presence of base, for example an inorganic base such as sodium carbonate or an organic base such as Hunig's base, in the presence of a suitable solvent such as DMF or THF at a temperature in the range of 0°C to reflux.

Compounds of formula (IXa) and (IXb) may be prepared by any of the procedures above for the preparation of compounds of formula (I), but wherein one of R' and R' is L.

Compounds of formula (X) are commercially available compounds, or they are known in the literature, or they are prepared by standard processes known in the art.

Process 7) and Process 8) and Process 10): Esters of formula (XIa), (XIb), (XIla), (XIId), (XVa) and (XVb) may be deprotected under standard conditions such as those described below, for example they may be deprotected with sodium hydroxide in methanol at room temperature.

Esters of formula (XIa), (XIb), (XIla), (XIId), (XVa) and (XVb) may be prepared by any of the procedures above for the preparation of compounds of formula (I), but wherein R¹¹ or R¹⁵ or R²⁶ is an ester.

It will be appreciated that certain of the various ring substituents in the compounds of the present invention may be introduced by standard aromatic substitution reactions or generated by conventional functional group modifications either prior to or immediately following the processes mentioned above, and as such are included in the process aspect of the invention. Such reactions and modifications include, for example, introduction of a substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution reactions include the introduction of a nitro group using concentrated nitric acid, the introduction of an acyl group using, for example, an acyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the introduction of a halogeno group. Particular examples of modifications include the reduction of a nitro group to an amino group by for example, catalytic hydrogenation with a nickel catalyst or treatment with iron in the presence of hydrochloric acid with heating; oxidation of alkylthio to alkylsulphinyl or alkylsulphonyl.

It will also be appreciated that in some of the reactions mentioned herein it may be necessary/desirable to protect any sensitive groups in the compounds. The instances where protection is necessary or desirable and suitable methods for protection are known to those skilled in the art. Conventional protecting groups may be used in accordance with standard practice (for illustration see T. W. Green, *Protective Groups in Organic Synthesis*, John Wiley and Sons, 1999). Thus, if reactants include groups such as amino, carboxy or hydroxy it may be desirable to protect the group in some of the reactions mentioned herein.

A suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxyacarbonyl group, for example a methoxycarbonyl, ethoxycarbonyl or *t*-butoxycarbonyl group, an arylmethoxycarbonyl group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxyacarbonyl group or an aroyl group may be removed for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *t*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulphuric or phosphoric acid or trifluoroacetic acid and an arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example dimethylaminopropylamine, or with hydrazine.

A suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethynyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an arylmethynyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

A suitable protecting group for a carboxy group is, for example, an esterifying group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a *t*-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

The protecting groups may be removed at any convenient stage in the synthesis using conventional techniques well known in the chemical art.

As stated hereinbefore the compounds defined in the present invention possess IBAT inhibitory activity. These properties may be assessed, for example, using an *in vitro* test assay

for studying the effect on bile acid uptake in IBAT-transfected cells (Smith L., Price-Jones M. J., Hughes K. T. and Jones N. R. A., *J Biomolecular Screening*, 3, 227-230) or *in vivo* by studying the effect on radiolabelled bile acid absorption in mice/rats (Lewis M. C., Brieady L. E. and Root C., *J Lip Res* 1995, 36, 1098-1105).

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore in association with a pharmaceutically-acceptable diluent or carrier.

The composition may be in a form suitable for oral administration, for example as a tablet or capsule, for parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion) as a sterile solution, suspension or emulsion, for topical administration as an ointment or cream or for rectal administration as a suppository.

In general the above compositions may be prepared in a conventional manner using conventional excipients.

The compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, will normally be administered to a warm-blooded animal at a unit dose within the range 5-5000 mg per square meter body area of the animal, i.e. approximately 0.02-100 mg/kg, preferably 0.02-50 mg/kg, and this normally provides a therapeutically-effective dose. A unit dose form such as a tablet or capsule will usually contain, for example 1-250 mg of active ingredient. Preferably a daily dose in the range of 1-50 mg/kg, particularly 0.1-10 mg/kg is employed. However the daily dose will necessarily be varied depending upon the host treated, the particular route of administration, and the severity of the illness being treated. Accordingly the optimum dosage may be determined by the practitioner who is treating any particular patient.

According to a further aspect of the present invention there is provided a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore for use in a method of prophylactic or therapeutic treatment of a warm-blooded animal, such as man.

We have found that the compounds defined in the present invention, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, are effective IBAT inhibitors, and accordingly have value in the treatment of disease states associated with hyperlipidaemic conditions.

Thus according to this aspect of the invention there is provided a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore for use as a medicament.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore, in the production of an IBAT inhibitory effect in a warm-blooded animal, such as man.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore, in the treatment of hyperlipidaemic conditions in a warm-blooded animal, such as man.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore in the manufacture of a medicament for use in the production of an IBAT inhibitory effect in a warm-blooded animal, such as man.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore in the manufacture of a medicament for use in the treatment of hyperlipidaemic conditions in a warm-blooded animal, such as man.

Herein, where "the production of an IBAT inhibitory effect" or "producing an IBAT inhibitory effect" is referred to particularly this refers to the treatment of hyperlipidaemic conditions. In another aspect, "the production of an IBAT inhibitory effect" or "producing an IBAT inhibitory effect" refers to the treatment of dyslipidemic conditions and disorders such as hyperlipidaemia, hypertriglyceridemia, hyperbetalipoproteinemia (high LDL), hyperprebetalipoproteinemia (high VLDL), hyperchylomicronemia, hypolipoproteinemia, hypercholesterolemia, hyperlipoproteinemia and hypoalphalipoproteinemia (low HDL). In another aspect "the production of an IBAT inhibitory effect" or "producing an IBAT inhibitory effect" refers to the treatment of different clinical conditions such as

atherosclerosis, arteriosclerosis, arrhythmia, hyper-thrombotic conditions, vascular dysfunction, endothelial dysfunction, heart failure, coronary heart diseases, cardiovascular diseases, myocardial infarction, angina pectoris, peripheral vascular diseases, inflammation of cardiovascular tissues such as heart, valves, vasculature, arteries and veins, aneurisms,

stenosis, restenosis, vascular plaques, vascular fatty streaks, leukocytes, monocytes and/or macrophage infiltration, intimal thickening, medial thinning, infectious and surgical trauma and vascular thrombosis, stroke and transient ischaemic attacks. In another aspect "the production of an IBAT inhibitory effect" or "producing an IBAT inhibitory effect" refers to the treatment of atherosclerosis, coronary heart diseases, myocardial infarction, angina pectoris, peripheral vascular diseases, stroke and transient ischaemic attacks in a warm-blooded animal, such as man.

According to a further feature of this aspect of the invention there is provided a method for producing an IBAT inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

According to a further feature of this aspect of the invention there is provided a method of treating hyperlipidemic conditions in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

The size of the dose required for the therapeutic or prophylactic treatment will necessarily be varied depending on the host treated, the route of administration and the severity of the illness being treated. A unit dose in the range, for example, 0.1-50 mg/kg preferably 0.1-10 mg/kg is envisaged.

The IBAT inhibitory activity defined hereinbefore may be applied as a sole therapy or may involve, in addition to a compound of the invention, one or more other substances and/or treatments. Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment. According to this aspect of the invention there is provided a pharmaceutical product comprising a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore and an additional IBAT inhibitory substance as defined hereinbefore and an additional hypolipidaemic agent for the conjoint treatment of hyperlipidaemia.

In another aspect of the invention, the compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, may be administered in

association with an HMG Co-A reductase inhibitor, or pharmaceutically acceptable salts, solvates, solvates of such salts or prodrugs thereof. Suitable HMG Co-A reductase inhibitors, pharmaceutically acceptable salts, solvates, solvates of such salts or prodrugs thereof are statins well known in the art. Particular statins are fluvastatin, lovastatin, pravastatin, simvastatin, atorvastatin, cerivastatin, beryastatin, delvastatin, mevastatin and (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulphonyl)amino]pyrimidin-5-yl](3R,5S)-3,5-dihydroxyhept-6-enoic acid (rosuvastatin), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof. A particular statin is atorvastatin, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof. A more particular statin is atorvastatin calcium salt. A further particular statin is (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulphonyl)amino]pyrimidin-5-yl](3R,5S)-3,5-dihydroxyhept-6-enoic acid, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof. A more particular statin is rosuvastatin calcium salt.

In an additional aspect of the invention, the compound of formula (I), or a

pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof may be administered in association with an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and/or a bile acid binder thereby avoiding a possible risk of excess of bile acids in colon caused by the inhibition of the ileal bile acid transport system. An excess of bile acids in the visceral contents may cause diarrhoea. Thus, the present invention also provides a treatment of a possible side effect such as diarrhoea in patients during therapy comprising the compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

An HMG Co-A-reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof will by its action decrease the endogenous cholesterol available for the bile acid synthesis and have an additive effect in combination with the compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof on lipid lowering.

Suitable bile acid binders for such a combination therapy are resins, such as cholestyramine and cholestipol. One advantage is that the dose of bile acid binder might be kept lower than the therapeutic dose for treatment of cholesteraemia in single treatment comprising solely a bile acid binder. By a low dose of bile acid binder any possible side effects caused by poor tolerance of the patient to the therapeutic dose could also be avoided.

Therefore in an additional feature of the invention, there is provided a method for producing an IBAT inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

Therefore in an additional feature of the invention, there is provided a method for producing an IBAT inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with a bile acid binder.

Therefore in an additional feature of the invention, there is provided a method for producing an IBAT inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in simultaneous, sequential or separate administration with a bile acid binder.

Therefore in an additional feature of the invention, there is provided a method of treating hyperlipidemic conditions in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

Therefore in an additional feature of the invention, there is provided a method of treating hyperlipidemic conditions in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a

prodrug thereof in simultaneous, sequential or separate administration with an effective amount of a bile acid binder.

Therefore in an additional feature of the invention, there is provided a method of treating hyperlipidemic conditions in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in simultaneous, sequential or separate administration with a bile acid binder.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in association with a pharmaceutically acceptable diluent or carrier.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a bile acid binder, in association with a pharmaceutically acceptable diluent or carrier.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a bile acid binder in association with a pharmaceutically acceptable diluent or carrier.

According to a further aspect of the present invention there is provided a kit

comprising a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

According to a further aspect of the present invention there is provided a kit comprising a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a bile acid binder.

According to a further aspect of the present invention there is provided a kit

comprising a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof and a bile acid binder.

According to a further aspect of the present invention there is provided a kit

comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a first unit dosage form;
- b) an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a second unit dosage form; and
- c) container means for containing said first and second dosage forms.

According to a further aspect of the present invention there is provided a kit

comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a first unit dosage form;
- b) a bile acid binder, in a second unit dosage form; and
- c) container means for containing said first and second dosage forms.

According to a further aspect of the present invention there is provided a kit

comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a first unit dosage form;
- b) an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a second unit dosage form;
- c) a bile acid binder, in a third unit dosage form; and

d) container means for containing said first, second and third dosage forms.

According to a further aspect of the present invention there is provided a kit

comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, together with a pharmaceutically acceptable diluent or carrier, in a first unit dosage form;

- b) an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a second unit dosage form; and
- c) container means for containing said first and second dosage forms.

According to a further aspect of the present invention there is provided a kit

comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, together with a pharmaceutically acceptable diluent or carrier, in a first unit dosage form;
- b) a bile acid binder, in a second unit dosage form; and
- c) container means for containing said first and second dosage forms.

According to a further aspect of the present invention there is provided a kit

comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, together with a pharmaceutically acceptable diluent or carrier, in a first unit dosage form;
- b) an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a second unit dosage form; and
- c) a bile acid binder, in a third unit dosage form; and
- d) container means for containing said first, second and third dosage forms.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in the manufacture of a medicament for use in the production of an IBAT inhibitory effect in a warm-blooded animal, such as man.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a bile acid binder, in the manufacture of a medicament for use in the production of an IBAT inhibitory effect in a warm-blooded animal, such as man.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a bile acid binder, in the manufacture of such a salt or a prodrug thereof, and a bile acid binder, in the manufacture

of a medicament for use in the production of an IBAT inhibitory effect in a warm-blooded animal, such as man.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in the manufacture of a medicament for use in the treatment of hyperlipidaemic conditions in a warm-blooded animal, such as man.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, a bile acid binder, in the manufacture of a medicament for use in the treatment of hyperlipidaemic conditions in a warm-blooded animal, such as man.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a bile acid binder, in the manufacture of a medicament for use in the treatment of hyperlipidaemic conditions in a warm-blooded animal, such as man.

According to a further aspect of the present invention there is provided a combination treatment comprising the administration of an effective amount of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier, with the simultaneous, sequential or separate administration of an effective amount of an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier to a warm-blooded animal, such as man in need of such therapeutic treatment.

According to a further aspect of the present invention there is provided a combination treatment comprising the administration of an effective amount of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier, with the simultaneous, sequential or separate administration of an effective amount of a bile acid binder, optionally together with a pharmaceutically acceptable diluent or carrier to a warm-blooded animal, such as man in need of such therapeutic treatment.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier, with the simultaneous, sequential or separate administration of an effective amount of an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in the manufacture of a medicament for use in the treatment of hyperlipidaemic conditions in a warm-blooded animal, such as man.

According to a further aspect of the present invention there is provided a combination treatment comprising the administration of an effective amount of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier, with the simultaneous, sequential or separate administration of an effective amount of an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable excipient, with the simultaneous, sequential or separate administration of an effective amount of a bile acid binder, optionally together with a pharmaceutically acceptable diluent or carrier to a warm-blooded animal, such as man in need of such therapeutic treatment.

According to an additional further aspect of the present invention there is provided a combination treatment comprising the administration of an effective amount of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier, with the simultaneous, sequential or separate administration one or more of the following agents selected from:

- a CETP (cholesteryl ester transfer protein) inhibitor, for example those referenced and described in WO 00/38725, page 7 line 22 - page 10, line 17 which are incorporated herein by reference;
- a cholesterol absorption antagonist for example azetidinones such as SCH 58235 and those described in US 5,767,115 which are incorporated herein by reference;
- a MTP (microsomal transfer protein) inhibitor for example those described in Science, 282, 751-54, 1998 which are incorporated herein by reference;
- a fibric acid derivative; for example clofibrate, gemfibrozil, fenofibrate, ciprofibrate and bezafibrate;
- a nicotinic acid derivative, for example, nicotinic acid (niacin), acipimox and niceritol;
- a phytosterol compound for example stanols;
- probucol;
- an anti-obesity compound for example orlistat (EP 129,748) and sibutramine (GB 2,184,122 and US 4,929,629);
- an antihypertensive compound for example an angiotensin converting enzyme inhibitor, an angiotensin II receptor antagonist, an adrenergic blocker, an alpha

and/or a beta adrenergic blocker, a mixed alpha/beta adrenergic blocker, an adrenergic stimulant, calcium channel blocker, a diuretic or a vasodilator;

> insulin;

> sulphonylureas including glibenclamide, tolbutamide;

> metformin; and/or

> acetose;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier to a warm-blooded animal, such as man in need of such therapeutic treatment.

Particular ACE inhibitors or pharmaceutically acceptable salts, solvates, solvate of such salts or a prodrugs thereof, including active metabolites, which can be used in combination with a compound of formula (I) include but are not limited to, the following compounds: alacepril, alatriptol, alitriptol calcium, anocovenin, benazepril, benazepril hydrochloride, benazeprilat, benzoylcaptopril, captopril, captopril-cysteine, captopril-glythothione, ceranapril, ceranopril, cilazapril, cilazaprilat, delapril, delapril-diacid, enalapril, enalaprilat, enapril, epiceptopril, fexoxythidine, fosfenopril, fosfenopril sodium, fosinopril sodium, fosinoprilat, fosinoprilic acid, glycypiril, hemophin-4, idapril, imidapril, indolapril, indolaprilat, ibenzapril, lisinopril, lyciumin A, lyciumin B, mixapril, moexipril, moexiprilat, murexacin A, murexacin B, murexacin C, pentopril, perindopril, perindoprilat, pivalopril, pivalopril, quinapril, quinapril hydrochloride, quinaprilat, ramipril, ramiprilat, spirapril, spirapril hydrochloride, spiraprilat, spirapril, spirapril hydrochloride, temocapril, temocapril hydrochloride, teprotide,trandolapril,trandolaprilat, uribapril, zalcicipril, zalciciprilat, zofenopril and zofenoprilat. Preferred ACE inhibitors for use in the present invention are ramipril, ramiprilat, lisinopril, enalapril and enalaprilat. More preferred ACE inhibitors for uses in the present invention are ramipril and ramiprilat.

Preferred angiotensin II antagonists, pharmaceutically acceptable salts, solvates, solvate of such salts or a prodrugs thereof for use in combination with a compound of formula (I) include, but are not limited to, compounds: candesartan, candesartan cilexetil, losartan, valsartan, irbesartan, tasosartan, telmisartan and eprosartan. Particularly preferred angiotensin II antagonists or pharmaceutically acceptable derivatives thereof for use in the present invention are candesartan and candesartan cilexetil.

In another aspect of the invention, the compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, may be administered in association with a PPAR alpha and/or gamma agonist, or pharmaceutically acceptable salts, solvates, solvates of such salts or prodrugs thereof. Suitable PPAR alpha and/or gamma agonists, pharmaceutically acceptable salts, solvates, solvates of such salts or prodrugs thereof are well known in the art. These include the compounds described in WO 01/12187, WO 01/12612, WO 99/62870, WO 99/62872, WO 99/62871, WO 98/57941, WO 01/40170, J Med Chem, 1996, 39, 665, Expert Opinion on Therapeutic Patents, 10 (5), 623-634 (in particular Chem, 1996, 39, 665, Expert Opinion on Therapeutic Patents, 10 (5), 623-634) and J Med Chem, the compounds described in the patent applications listed on page 634) and J Med Chem, 2000, 43, 527 which are all incorporated herein by reference. Particularly a PPAR alpha and/or gamma agonist refers to WY-14643, clofibrate, fenofibrate, bezafibrate, GW 9578, troglitazone, pioglitazone, rosiglitazone, ciglitazone, proglitazone, BRI-49634, KRP-297, JTT-501, SB 213068, GW 1929, GW 7845, GW 0207, L-796449, L-165041, NN622/Ragaglitazar, BMS 298585 and GW 2433. Particularly a PPAR alpha and/or gamma agonist refers to (S)-2-ethoxy-3-(4-(2-(4-methanesulphonyloxyphenyl)ethoxy)phenyl)propanoic acid and pharmaceutically acceptable salts thereof.

Therefore in an additional feature of the invention, there is provided a method for producing an IBAT inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

Therefore in an additional feature of the invention, there is provided a method of treating hyperlipidemic conditions in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable

salt, solvate, solvate of such a salt or a prodrug thereof, and a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in association with a pharmaceutically acceptable diluent or carrier.

According to a further aspect of the present invention there is provided a kit

- 5 comprising a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

According to a further aspect of the present invention there is provided a kit

comprising:

- 10 a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a first unit dosage form;
 b) a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a second unit dosage form; and
 c) container means for containing said first and second dosage forms.

15 According to a further aspect of the present invention there is provided a kit

comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, together with a pharmaceutically acceptable diluent or carrier, in a first unit dosage form;

- 20 b) a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a second unit dosage form; and
 c) container means for containing said first and second dosage forms.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in the manufacture of a medicament for use in the production of an IBAT inhibitory effect in a warm-blooded animal, such as man.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in the manufacture of a medicament for use in the treatment of hyperlipidaemic conditions in a warm-blooded animal, such as man.

According to a further aspect of the present invention there is provided a combination treatment comprising the administration of an effective amount of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier, with the

- 5 simultaneous, sequential or separate administration of an effective amount of a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier to a warm-blooded animal, such as man in need of such therapeutic treatment.

In addition to their use in therapeutic medicine, the compounds of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, are also useful as pharmacological tools in the development and standardisation of *in vitro* and *in vivo* test systems for the evaluation of the effects of inhibitors of IBAT in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of the search for new therapeutic agents.

Many of the intermediates described herein are novel and are thus provided as a

- 15 further feature of the invention. For Example compounds of formula (XIa), (XIb), (XIIa), (XIIb), (XVa) and (XVb) show IBAT inhibitory activity when tested in the above referenced *in vitro* test assay and are thus claimed as a further feature of the invention.

Thus in a further feature of the invention, there is provided a compound of formula (XIa), (XIb), (XIIa), (XIIb), (XVa) or (XVb), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

Therefore according to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (XIa), (XIb), (XIIa), (XIIb), (XVa) or (XVb), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore in association with a pharmaceutically-acceptable diluent or carrier.

According to an additional aspect of the present invention there is provided a compound of the formula (XIa), (XIb), (XIIa), (XIIb), (XVa) or (XVb), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore for use in a method of prophylactic or therapeutic treatment of a warm-blooded animal, such as man.

Thus according to this aspect of the invention there is provided a compound of the formula (XIa), (XIb), (XIIa), (XIIb), (XVa) or (XVb), or a pharmaceutically acceptable salt,

solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore for use as a medicament.

According to another feature of the invention there is provided the use of a compound of the formula (XIa), (XIb), (XIIfa), (XIIfb), (XV'a) or (XV'b), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof as defined hereinbefore in the manufacture of a medicament for use in the production of an IBAT inhibitory effect in a warm-blooded animal, such as man.

According to another feature of the invention there is provided the use of a compound of the formula (XIa), (XIb), (XIIfa), (XIIfb), (XV'a) or (XV'b), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof as defined hereinbefore in the manufacture of a medicament for use in the treatment of hyperlipidaemic conditions in a warm-blooded animal, such as man.

According to a further feature of this aspect of the invention there is provided a method for producing an IBAT inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (XIa), (XIb), (XIIfa), (XIIfb), (XV'a) or (XV'b), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

According to a further feature of this aspect of the invention there is provided a method of treating hyperlipidaemic conditions in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (XIa), (XIb), (XIIfa), (XIIfb), (XV'a) or (XV'b), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

In the above other pharmaceutical composition, process, method, use and medicament manufacture features, the alternative and preferred embodiments of the compounds of the invention described herein also apply.

Examples

The invention will now be illustrated in the following non limiting examples, in which standard techniques known to the skilled chemist and techniques analogous to those described in these examples may be used where appropriate, and in which, unless otherwise stated:

(i) evaporations were carried out by rotary evaporation in vacuo and work up procedures were carried out after removal of residual solids such as drying agents by filtration;

(ii) all reactions were carried out under an inert atmosphere at ambient temperature, typically in the range 18-25°C, with solvents of HPLC grade under anhydrous conditions, unless otherwise stated;

(iii) column chromatography (by the flash procedure) was performed on Silica gel 40-63 µm (Merck);

(iv) yields are given for illustration only and are not necessarily the maximum attainable;

(v) the structures of the end products of the formula (I) were generally confirmed by nuclear (generally proton) magnetic resonance (NMR) and mass spectral techniques; magnetic resonance chemical shift values were measured in deuterated CD₃OD (unless otherwise stated) on the delta scale (ppm downfield from tetramethylsilane); proton data is quoted unless

otherwise stated; spectra were recorded on a Varian Mercury-300 MHz, Varian Unity plus-400 MHz, Varian Unity plus-600 MHz or on Varian Inova-500 MHz spectrometer; and peak multiplicities are shown as follows: s, singlet; d, doublet; dd, double doublet; t, triplet; tt, triple triplet; q, quartet; tq, triple quartet; m, multiplet; br, broad; LCMS were recorded on a

Waters ZMD, LC column xTerra MS C₁₈(Waters), detection with a HP 1100 MS-detector diode array equipped; mass spectra (MS) (loop) were recorded on VG Platform II (Fisons Instruments) with a HP-1100 MS-detector diode array equipped; unless otherwise stated the mass ion quoted is (MH)⁺;

(vi) unless further details are specified in the text, analytical high performance liquid chromatography (HPLC) was performed on Prep LC 2000 (Waters), Kromasil C₁₈ 7µm, (Akzo Nobel), MeCN and de-ionised water 100 mM ammonium acetate as mobile phases, with suitable composition;

(vii) intermediates were not generally fully characterised and purity was assessed by thin layer chromatography (TLC), HPLC, infra-red (IR), MS or NMR analysis;

(viii) where solutions were dried sodium sulphate was the drying agent;

(ix) where an "ISOLUTE" column is referred to, this means a column containing 2g of silica, the silica being contained in a 6 ml disposable syringe and supported by a porous disc of 54Å pore size, obtained from International Sorbent Technology under the name "ISOLUTE"; "ISOLUTE" is a registered trade mark;

(x) the following abbreviations may be used hereinbefore or hereinafter:-

DCM dichloromethane;
DMF N,N-dimethylformamide;

TFA trifluoroacetic acid;
 TBTU o-Benzotriazol-1-yl-N,N,N',N'-tetramethyluronium tetrafluoroborate;
 EtOAc ethyl acetate; and
 MeCN acetonitrile.

Example 1

1,1-Dioxo-3-(R)-3-butyl-3-ethyl-5-(R)-5-phenyl-8-[N-(R)- α -carboxybenzyl]carbamoylmethoxy]-2,3,4,5-tetrahydro-1,4-benzothiazepine; and
 1,1-Dioxo-3-(S)-3-butyl-3-ethyl-5-(S)-5-phenyl-8-[N-(R)- α -carboxybenzyl]carbamoylmethoxy]-2,3,4,5-tetrahydro-1,4-benzothiazepine

(+)-trans-1,1-Dioxo-3-butyl-3-ethyl-5-phenyl-8-(carboxymethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine (Method 1; 13 mg, 0.03 mmol) methyl (2R)-amino(phenyl)acetate (7.5 mg, 0.037 mmol) and diisopropylethylamine (24 mg, 0.19 mmol) were dissolved in DCM (1.5 ml). The mixture was stirred for 10 min and then TBTU (12 mg, 0.037 mmol) was added and the reaction mixture was stirred for 30 min. The solvent was removed under reduced pressure. The residue was dissolved in ethanol (2 ml) and sodium hydroxide (2 mg) was added. The mixture was stirred for 30 min and the solvent was evaporated. The residue was purified by chromatography (DCM : EtOAc : AcOH, 100:10:3) giving the title compound (5.5 mg, 32%).
 M/z: 565.3 (MH⁺), 563.2 (M⁺).

Example 2

1,1-Dioxo-3-(R)-3-butyl-3-ethyl-5-(R)-5-phenyl-8-[N-(R)- α -[N-(carboxymethyl)carbamoyl]benzyl]carbamoylmethoxy]-2,3,4,5-tetrahydro-1,4-benzothiazepine; and
 1,1-Dioxo-3-(S)-3-butyl-3-ethyl-5-(S)-5-phenyl-8-[N-(R)- α -[N-(carboxymethyl)carbamoyl]benzyl]carbamoylmethoxy]-2,3,4,5-tetrahydro-1,4-benzothiazepine

An equal mixture of 1,1-dioxo-3-(R)-3-butyl-3-ethyl-5-(R)-5-phenyl-8-[N-(R)- α -[N-(t-butoxycarbonylmethyl)carbamoyl]benzyl]carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine and 1,1-dioxo-3-(S)-3-butyl-3-ethyl-5-(S)-5-phenyl-8-[N-(R)- α -[N-(t-butoxycarbonylmethyl)carbamoyl]benzyl]carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine (Method 2; 27 mg, 0.040 mmol) were dissolved in 2 ml DCM.

Trifluoroacetic acid (0.2 ml, 2.60 mmol) was added and the mixture was stirred overnight at ambient temperature. The reaction mixture was concentrated under reduced pressure and then

purified with preparative HPLC using an MeCN/ammonium acetate buffer gradient (5/95 to 60/40) as eluent. The MeCN was evaporated and lyophilisation of the remaining solution resulted in the title products in 69% yield (16 mg). NMR (400 MHz, MeOD): 0.81 (t, 3H), 0.89 (t, 3H), 1.11-1.35 (m, 4H), 1.41-1.50 (m, 1H), 1.52-1.62 (m, 1H), 1.74-1.84 (m, 1H), 2.17-2.28 (m, 1H), 3.34 (ABq, 2H), 3.87 (ABq, 2H), 4.63-4.66 (m, 2H), 5.61 (s, 1H), 6.00 (s, 1H), 6.59-6.64 (m, 1H), 6.95-7.01 (m, 1H), 7.27-7.44 (m, 10H), 7.64-7.67 (m, 1H); m/z: 622 (M+1).

Example 3

0 3,5-trans-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-7-bromo-8-[N-(R)- α -[N-(carboxymethyl)carbamoyl]benzyl]carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine
 3,5-trans-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-7-bromo-8-(carboxymethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine (Method 5; 50 mg, 0.10 mmol) was dissolved in DCM (3 ml).
 5 Lutidine (0.022 ml, 0.198 mmol), TBTU (38 mg, 0.118 mmol) and (R)- α -[N-(t-butoxycarbonylmethyl)carbamoyl]benzylamine (Method 4; 44 mg, 0.167 mmol) were added successively. The mixture was stirred over night at ambient temperature. The solution was concentrated to 1 ml and TFA (1.3 ml) was added. The mixture was concentrated after 1.5h and the residue was purified using preparative HPLC. A gradient from 40% to 60% of MeCN in 0.1 M ammonium acetate buffer was used as eluent. Lyophilisation yielded 39 mg (57%).
 10 NMR (400 MHz) 0.75 (t, 3H), 0.95 (t, 3H), 1.2-1.4 (m, 6H), 1.75-1.9 (m, 1H), 2.2-2.4 (m, 1H), 3.35 (dd, 2H), 3.85 (dd, 2H), 4.7-4.8 (m, 2H), 5.6 (s, 1H), 6.0 (d, 1H), 6.8 (d, 1H), 7.25-7.5 (m, 10H), 7.6 (d, 1H); m/z: 700 (M) and 702 (M+2)⁺.

Example 4

25 3,5-trans-1,1-Dioxo-3-(S)-3-ethyl-3-butyl-4-hydroxy-5-(S)-5-phenyl-7-bromo-8-[N-(R)- α -[N-(carboxymethyl)carbamoyl]benzyl]carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine
 3,5-trans-1,1-Dioxo-3-(R)-3-ethyl-3-butyl-4-hydroxy-5-(R)-5-phenyl-7-bromo-8-[N-(R)- α -[N-(carboxymethyl)carbamoyl]benzyl]carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine

3,5-*trans*-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-7-bromo-8-(*N*-(*R*)- α -[*N*-(carboxymethyl)carbamoyl]benzyl)carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine (Example 3; 14 mg, 0.02 mmol) was dissolved in 2 ml DCM. *m*-Chloroperoxybenzoic acid (5.5 mg, 0.022 mmol) was added and the mixture was stirred for 30 min. The diastereomers of the title compound were separated using preparative HPLC on a C8 column. A gradient from 30% to 60% of MeCN in 0.1 M ammonium acetate buffer was used as eluent. The two compounds were lyophilized and the first eluting diastereomer was obtained in 5.4 mg and the second in 4.9 mg. *M/z*: 716 (M) and 718 (M+2)⁺. NMR (400 MHz) (diastereomer 1) 0.86 (t, 3H), 0.95 (t, 3H), 1.1-1.4 (m, 3H), 1.4-1.55 (m, 2H), 1.68-1.8 (m, 1H), 2.0-2.2 (m, 2H), 3.4 (dd, 2H), 3.88 (Abq, 2H), 4.76 (Abq, 2H), 5.6 (s, 1H), 6.45 (s, 1H), 6.88 (s, 1H), 7.25-7.50 (m, 10H), 7.56 (s, 1H). NMR (diastereomer 2) (400 MHz) 0.87 (t, 3H), 0.95 (t, 3H), 1.1-1.4 (m, 3H), 1.4-1.55 (m, 2H), 1.68-1.8 (m, 1H), 2.0-2.22 (m, 2H), 3.4 (dd, 2H), 3.82 (Abq, 2H), 4.76 (Abq, 2H), 5.6 (s, 1H), 6.46 (s, 1H), 6.88 (s, 1H), 7.25-7.50 (m, 10H), 7.57 (s, 1H).

Example 5

3,5-*trans*-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-(*N*-(*R*)- α -[*N*-(carboxymethyl)carbamoyl]benzyl)carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine

3,5-*trans*-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-(carboxymethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine (Method 6; 50 mg, 0.105 mmol) was dissolved in DCM (2 ml). 2,6-Lutidine (0.025 ml, 0.215 mmol), TBTU (45 mg, 0.140 mmol) and (*R*)- α -[*N*-(4-butoxycarbonylmethyl)carbamoyl]benzylamine (Method 4; 43 mg, 0.163 mmol) were added successively. The mixture was stirred for 2 hours at ambient temperature. The solution was concentrated and the intermediate ester was purified by chromatography on silica using DCM/EtOAc (9/1) as eluent. The solvent was evaporated to yield 45 mg (60%) *M/z*: 724. The ester was dissolved in 3 ml DCM and hydrolysed by addition of TFA (1 ml). After 2 hours the mixture was concentrated and purified using preparative HPLC. A gradient of MeCN from 40% to 60% in 0.1 M ammonium acetate buffer was used as eluent.

Lyophilisation yielded 33 mg (80%). NMR (400 MHz): 0.75-0.85 (m, 3H), 0.85-0.95 (m, 3H), 1.1-1.65 (m, 6H), 1.75-1.9 (m, 1H), 2.0 (s, 3H), 2.2-2.4 (m, 1H), 3.1-3.55 (m, 2H), 3.85 (Abq,

2H), 4.6-4.8 (m, 2H), 5.6 (s, 1H), 5.98-6.03 (m, 1H), 6.4 (s, 1H), 7.25-7.56 (m, 11H); *m/z*: 668.

Example 6

3,5-*trans*-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-(*N*-(*R*)- α -[*N*-(2-sulphoethyl)carbamoyl]-4-hydroxybenzyl)carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine ammonium salt

3,5-*trans*-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-(carboxymethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine (Method 6; 33 mg, 0.070 mmol) was dissolved in DMF (3 ml). 2-((2*R*)-2-Amino-2-(4-hydroxyphenyl)ethanol)amino)ethanesulphonic acid (Method 8; 23 mg, 0.084 mmol), *N*-methylmorpholine (0.025 ml, 0.227 mmol) and TBTU (27 mg, 0.084 mmol) were added successively and the mixture was stirred overnight. The solvent was removed and the crude product was purified using preparative HPLC. A gradient from 40% to 70% of MeCN in 0.1 M ammonium acetate buffer was used as eluent. Lyophilisation yielded 42 mg (80%) of the ammonium salt. NMR (400 MHz): 0.73-0.85 (m, 3H), 0.85-0.98 (m, 3H), 1.1-1.7 (m, 6H), 1.75-1.9 (m, 1H), 2.0 (s, 3H), 2.15-2.4 (m, 1H), 2.85-3.0 (m, 2H), 3.1-3.55 (m, 2H), 3.5-3.65 (m, 2H), 4.6-4.8 (m, 2H), 5.35-5.39 (m, 1H), 5.98-6.05 (m, 1H), 6.4 (s, 1H), 6.75 (d, 2H), 7.15-7.5 (m, 8H); *m/z*: 734.

Example 7

1,1-Dioxo-3-(*S*)-3-ethyl-3-butyl-5-(*S*)-5-phenyl-7-methylthio-8-(*N*-(*R*)- α -[*N*-(carboxymethyl)carbamoyl]benzyl)carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine diethylamine salt

1,1-Dioxo-3-(*R*)-3-ethyl-3-butyl-5-(*R*)-5-phenyl-7-methylthio-8-(*N*-(*R*)- α -[*N*-(carboxymethyl)carbamoyl]benzyl)carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine diethylamine salt

The diastereomeric mixture of 3,5-*trans*-1,1-dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-(*N*-(*R*)- α -[*N*-(carboxymethyl)carbamoyl]benzyl)carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine (Example 5; 17 mg, 0.026 mmol) was separated by chiral chromatography on a Chirobiotic V chiral stationary phase. Two columns (250 x 20 mm) in series were used. A mobile phase consisting of 80% MeOH in water with 0.1% Et₃N and 0.1% HOAc was used as eluent. The first eluting diastereomer was collected in a 50 ml

fraction and the solvent was removed under reduced pressure. Et₃N remained according to NMR-analysis and the diastereomer was purified by chromatography over 0.5 g SiO₂ using DCM/MeOH (9/1) as eluent. The solvent was removed and the product was dissolved in water and some MeCN. Lyophilisation yielded a white solid, which was dissolved in MeOH and filtered. A second lyophilisation yielded the diastereomer as the Et₃N salt in 1 mg (4%). M/z: 668. NMR (HOAc-d₄) was consistent with Example 5. The e.e. was determined as 99%. The second eluting diastereomer was collected in a 200 ml fraction and the solvent was removed under reduced pressure. The residue was purified using preparative HPLC on a C8 column. A gradient from 35% to 50% MeCN in 0.1 M ammonium acetate was used as eluent.

10 Lyophilisation yielded the diastereomer as the Et₃N salt in 3 mg (17 mg). M/z: 668. The e.e. was determined as 97%.

Preparation of Starting Materials

The starting materials for the Examples above are either commercially available or are readily prepared by standard Methods from known materials. For Example, the following reactions are an illustration, but not a limitation, of some of the starting materials used in the above reactions.

Method 1

20 (+)-trans-1,1-Dioxo-3-butyl-3-butyl-3-ethyl-5-phenyl-8-(carboxymethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine

1,1-Dioxo-3-butyl-3-ethyl-5-phenyl-8-hydroxy-2,3,4,5-tetrahydro-1,4-benzothiazepine (prepared according to WO/9605188; 83 mg, 0.22 mmol), ethyl bromoacetate (55 mg, 0.33 mmol) and sodium carbonate (70 mg, 0.66 mmol) in acetonitrile (3 ml) were warmed to reflux for 40 hours. The solvent was removed under reduced pressure and the crude product was dissolved in ethanol (4 ml). Sodium hydroxide (0.1 g) was added and the mixture was warmed to reflux for 1 hour. The solvent was removed under reduced pressure and the residue was partitioned between DCM and 2 M acetic acid. The organic layer was dried over sodium sulphate and the solvent was removed under reduced pressure. The residue was purified by chromatography (HOAc: formic acid, 500:1) to give 61 mg (64%) of the title compound.

30 NMR (500 MHz, CDCl₃): 0.86 (t, 3H), 0.92 (t, 3H), 1.0-1.05 (m, 1H), 1.2-1.4 (m, 3H), 1.6-1.75 (m, 2H), 1.85-1.95 (m, 1H), 2.38-2.47 (m, 1H), 3.45 (s, 2H), 4.5 (s, 2H), 6.17 (s, 1H), 6.75 (d, 1H), 6.86 (dd, 1H), 7.37-7.5 (m, 5H), 7.64 (d, 1H).

Method 2

1,1-Dioxo-3(R)-3-butyl-3-ethyl-5-(R)-5-phenyl-8-(N-(R)-α-[N-(t-butoxycarbonylmethyl)carbamoyl]benzyl)carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine; and

1,1-Dioxo-3(S)-3-butyl-3-ethyl-5-(S)-5-phenyl-8-(N-(R)-α-[N-(t-butoxycarbonylmethyl)carbamoyl]benzyl)carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine

5 (+)-trans-1,1-Dioxo-3-butyl-3-ethyl-5-phenyl-8-(carboxymethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine (Method 1; 17.5 mg, 0.041 mmol) was dissolved in DCM (3 ml), 2,6-Lutidine (0.010 ml, 0.086 mmol), TBUTU (16.4 mg, 0.051 mmol) and (R)-α-[N-(t-butoxycarbonylmethyl)carbamoyl]benzylamine (Method 4; 16.3 mg, 0.062 mmol) were added successively. The mixture was stirred for 1 hour at ambient temperature. The solution was concentrated and the crude product was purified by chromatography on silica using DCM/EtOAc (8/2) as eluent. The solvent was evaporated and the title products were obtained in 98% yield (27 mg). M/z: 678 (M+1).

Method 3

(R)-N-Benzoyloxycarbonyl-α-[N-(t-butoxycarbonylmethyl)carbamoyl]benzylamine

(2R)-{[(Benzoyloxy)carbonyl]amino}(phenyl)acetic acid (10 g, 35.0 mmol) and t-butylglycine hydrochloride (6.3 g, 37.4 mmol) were dissolved in DCM (200 ml) with 2,6-lutidine (8.2 ml, 70.4 mmol). After stirring 5 min at 0°C TBUTU (12.4 g, 38.6 mmol) was added and stirring was continued for 1.5 hours at 0°C and 3.75 hours at room temperature. The reaction mixture was washed with water (2 x 100 ml), dried (MgSO₄) and purified with flash chromatography (DCM:EtOAc 7:1→5:1) to give the title compound (13 g, 94 %). NMR (500 MHz, CDCl₃): 1.45 (s, 9H), 3.84 (d, 1H), 4.00 (dd, 1H), 5.10 (m, 2H), 5.28 (brs, 1H), 6.13 (brs, 1H), 6.23 (brs, 1H), 7.30-7.44 (m, 10H).

Method 4

(R)-α-[N-(t-Butoxycarbonylmethyl)carbamoyl]benzylamine

(R)-N-Benzoyloxycarbonyl-α-[N-(t-butoxycarbonylmethyl)carbamoyl]benzylamine (Method 3; 12.8 g, 32.2 mmol) was dissolved in EtOH (99%, 200 ml) and toluene (50 ml). PUC (10%, 0.65 g) was added and hydrogenation was performed at atmospheric pressure for 5.5 hours at room temperature. The reaction mixture was filtered through diatomaceous earth

and the solvents were evaporated to give the title compound (3.4 g, 99 %). NMR (600 MHz, CDCl₃): 1.45 (s, 9H), 3.93 (m, 2H), 4.54 (s, 1H), 7.31-7.42 (m, 5H), 7.51 (m, 1H).

Method 5

3,5-*trans*-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-7-bromo-8-(carboxymethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine

The title compound was prepared as described in Method 6 starting from (+/-)-*trans*-7-bromo-3-butyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-8-ol (1,1-dioxide (WO96/05188; 81 mg, 0.18 mmol). The intermediate ethyl ester was obtained in 94% yield (m/z: 538(M)). The product was obtained in 50 mg (58%). NMR 0.75 (t, 3H), 0.95 (t, 3H), 1.2-1.45 (m, 6H), 1.75-1.9 (m, 1H), 2.2-2.4 (m, 1H), 3.35 (dd, 2H), 4.8 (s, 2H), 6.0 (s, 1H), 6.8 (s, 1H), 7.3-7.5 (m, 5H), 7.55 (s, 1H); m/z: 510 (M) and 512 (M+2)⁺.

Method 6

3,5-*trans*-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-(carboxymethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine

The title compound was prepared from 3,5-*trans*-1,1-dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-hydroxy-2,3,4,5-tetrahydro-1,4-benzothiazepine (Method 7; 153 mg, 0.36 mmol) using the procedure described in Method 1. The intermediate ethyl ester was extracted between diluted HCl and DCM. The DCM phase was washed with brine, dried with Na₂SO₄ and concentrated. M/z 506. The crude product was dissolved in THF/H₂O (3/1; 4 ml) and LiOH (22 mg, 0.91 mmol) was added. The mixture was stirred for 2h and the solvent was removed under reduced pressure. The crude product was purified using preparative HPLC. A gradient from 40% to 60% MeCN in 0.1 M ammonium acetate buffer was used as eluent. The MeCN was removed under reduced pressure and the remaining aqueous solution was acidified using 5% HCl and was then extracted with DCM. The DCM layer was dried with Na₂SO₄ and concentrated. The crude product was co-evaporated with diethyl ether. The obtained crystals were filtered off and dried. Mass: 158 mg (91%). NMR 0.75 (t, 3H), 0.9 (t, 3H), 1.1-1.7 (m, 6H), 1.7-1.9 (m, 1H), 2.0 (s, 3H), 2.2-2.4 (m, 1H), 3.3 (dd, 2H), 4.75 (s, 2H), 6.0 (s, 1H), 6.4 (s, 1H), 7.3-7.5 (m, 6H); m/z: 478.

Method 7

3,5-*trans*-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-hydroxy-2,3,4,5-tetrahydro-1,4-benzothiazepine

(+/-)-*trans*-7-Bromo-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide (prepared according to WO 96/05188; 300 mg, 0.64 mmol) was dissolved in 5 ml DMF under N₂/atmosphere. Sodium thiomethylate (150 mg, 2.14 mmol) was added and the mixture was heated to 110°C for 2h. The solvent was removed under reduced pressure and the residue was extracted between 5% HCl and EtOAc. The organic phase was washed with brine, dried with Na₂SO₄ and concentrated. The product was purified using preparative HPLC. A gradient from 40% to 100% of MeCN in 0.1 M ammonium acetate buffer was used as eluent. Lyophilisation yielded 153 mg, 57%. M/z: 420.


Method 8

2-((2R)-2-Amino-2-(4-hydroxyphenyl)ethanol)ethanesulphonic acid

N-Boc-(D)-4-hydroxyphenylglycine (1.00 g, 3.21 mmol) was dissolved in DMF (5 ml) and tetrabutylammonium taurine (2.36 g, 6.42 mmol) was added together with additional DMF (5 ml). The resulting suspension was cooled on ice and TBUTU (1.24 g, 3.85 mmol) was added. The ice bath was removed after 30 min and the mixture was stirred for 2 hours before it was filtered and concentrated. TFA in DCM (20%, 20 ml) was added and the reaction mixture was stirred over night. Ethanol (20 ml) was added and the solvents evaporated. The crude product was refluxed in ethanol (100 ml) for 1 hour. Filtration yielded the pure title compound as a white solid, 626 mg (71%). NMR (DMSO-d₆): 2.4-2.6 (m, 2H), 3.2-3.4 (m, 2H), 4.79 (s, 1H), 6.78 (d, 2H), 7.23 (d, 2H), 8.22 (t, 1H), 8.4 (m, 3H), 9.7 (s, 1H).

Claims

1. A compound of formula (I):



5

wherein:

One of R¹ and R² are selected from hydrogen, C₁₋₆alkenyl or C₂₋₆alkenyl and the other is selected from C₁₋₆alkyl or C₂₋₆alkenyl;

R³ is selected from hydrogen, hydroxy, C₁₋₆alkyl, C₁₋₆alkoxy and C₁₋₆alkoxyloxy;

10 R⁴ is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphonamoyl, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆alkoxy, C₁₋₆alkenyl, C₁₋₆alkoxyloxy,

N-(C₁₋₆alkyl)amino, N,N-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, N-(C₁₋₆alkyl)carbamoyl,


N,N-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkylS(O)_n, wherein n is 0 to 2, C₁₋₆alkoxy, carbonyl,

N-(C₁₋₆alkyl)sulphonamoyl and N,N-(C₁₋₆alkyl)₂sulphonamoyl;

15 N is 0-5;

wherein:

one of R⁴ and R⁵ is a group of formula (IA):



R² and R⁴ and the other of R⁴ and R⁵ are independently selected from hydrogen, halo,

20 nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphonamoyl, C₁₋₆alkyl,

C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆alkoxy, C₁₋₆alkenyl, C₁₋₆alkoxyloxy, N-(C₁₋₆alkyl)amino,

N,N-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, N-(C₁₋₆alkyl)carbamoyl,

N,N-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkylS(O)_n, wherein n is 0 to 2, C₁₋₆alkoxy, carbonyl,

N-(C₁₋₆alkyl)sulphonamoyl and N,N-(C₁₋₆alkyl)₂sulphonamoyl; wherein R³ and R⁶ and the other of R⁴ and R⁵ may be optionally substituted on carbon by one or more R¹⁶;

X is -O-, -N(R¹⁷), -S(O)_n or -CH(R¹⁸); wherein R¹⁷ is hydrogen or C₁₋₆alkyl and n is 0-2;

5 Ring A is aryl or heteroaryl; wherein Ring A is optionally substituted by one or more substituents selected from R¹⁷;


R⁷ is hydrogen, C₁₋₆alkyl, carbocyclyl or heterocyclyl; wherein R⁷ is optionally substituted by one or more substituents selected from R¹⁸;

R⁸ is hydrogen or C₁₋₆alkyl;

10 R⁹ is hydrogen or C₁₋₆alkyl;

R¹⁰ is hydrogen, C₁₋₆alkyl, carbocyclyl or heterocyclyl; wherein R¹⁰ is optionally substituted by one or more substituents selected from R¹⁹;

R¹¹ is carboxy, sulpho, sulphinio, phosphono, -P(O)(OR²⁰)(OR²¹), -P(O)(OH)(OR²²), -P(O)(OH)(R²³) or -P(O)(OR²⁴)(R²⁵) wherein R²⁰ and R²¹ are independently selected from C₁₋₆alkyl; or R¹¹ is a group of formula (IB):



wherein:


20 Y is -N(R²⁶), -N(R²⁷)C(O)-, -O-, and -S(O)_n; wherein n is 0-2 and R²⁶ is hydrogen or C₁₋₆alkyl;

R¹² is hydrogen or C₁₋₆alkyl;

R¹³ and R¹⁴ are independently selected from hydrogen, C₁₋₆alkyl, carbocyclyl or heterocyclyl; wherein R¹³ and R¹⁴ may be independently optionally substituted by one or more substituents selected from R²⁰;

25 R¹⁵ is carboxy, sulpho, sulphinio, phosphono, -P(O)(OR²⁸)(OR²⁹), -P(O)(OH)(OR³⁰), -P(O)(OH)(R³¹) or -P(O)(OR³²)(R³³) wherein R²⁸ and R²⁹ are independently selected from C₁₋₆alkyl; or R¹⁵ is a group of formula (IC):

- 54 -



wherein:

R^{24} is selected from hydrogen or C_{1-6} alkyl;

R^{25} is selected from hydrogen, C_{1-6} alkyl, carboxycyl, heterocycyl or R^{21} , wherein said C_{1-6} alkyl, carboxycyl or heterocycyl may be independently optionally substituted by one or more substituents selected from R^{23} ;

R^{26} is selected from carboxy, sulpho, sulphinio, phosphono, tetrazolyl,

$-P(O)(OR^9)(OR^9)$, $-P(O)(OH)(R^9)$ or $-P(O)(OR^9)(R^9)$ wherein R^8 and R^9 are independently selected from C_{1-6} alkyl;

p is 1-3; wherein the values of R^{15} may be the same or different;

q is 0-1;

r is 0-3; wherein the values of R^{14} may be the same or different;

m is 0-2; wherein the values of R^{10} may be the same or different;

n is 1-3; wherein the values of R^7 may be the same or different;

z is 0-3; wherein the values of R^{25} may be the same or different;

R^{16} , R^{17} and R^{18} are independently selected from halo, nitro, cyano, hydroxy, amino,

carboxy, carbamoyl, mercapto, sulphanoyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-6} alkynyl, C_{1-6} alkoxy,

C_{1-6} alkanoyl, C_{1-6} alkanyloxy, $N-(C_{1-6}alkyl)amino$, $N,N-(C_{1-6}alkyl)_2amino$, $C_{1-6}alkylS(O)_n$,

$C_{1-6}alkanylamino$, $N-(C_{1-6}alkyl)carbamoyl$, $N,N-(C_{1-6}alkyl)_2carbamoyl$, $C_{1-6}alkylS(O)_n$,

wherein a is 0 to 2, $C_{1-6}alkoxy$ carbonyl, $N-(C_{1-6}alkyl)sulphanoyl$ and

$N,N-(C_{1-6}alkyl)_2sulphanoyl$; wherein R^{16} , R^{17} and R^{18} may be independently optionally

substituted on carbon by one or more R^{21} ;

R^{19} , R^{20} and R^{22} are independently selected from halo, nitro, cyano, hydroxy,

amino, carboxy, carbamoyl, mercapto, sulphanoyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-6} alkynyl,

$C_{1-6}alkoxy$, $C_{1-6}alkanoyl$, $C_{1-6}alkanyloxy$, $N-(C_{1-6}alkyl)amino$, $N,N-(C_{1-6}alkyl)_2amino$,

$C_{1-6}alkanylamino$, $N-(C_{1-6}alkyl)carbamoyl$, $N,N-(C_{1-6}alkyl)_2carbamoyl$, $C_{1-6}alkylS(O)_n$,

wherein a is 0 to 2, $C_{1-6}alkoxy$ carbonyl, $N-(C_{1-6}alkyl)sulphanoyl$, $N,N-(C_{1-6}alkyl)_2sulphanoyl$,

carboxycyl, heterocycyl, sulpho, sulphinio, amidino, $(C_{1-6}alkyl)silyl$, phosphono,

$-P(O)(OR^9)(OR^9)$, $-P(O)(OH)(OR^9)$, $-P(O)(OH)(R^9)$ or $-P(O)(OR^9)(R^9)$ wherein R^8 and R^9 are

- 55 -

independently selected from C_{1-6} alkyl; wherein R^{19} and R^{20} may be independently optionally substituted on carbon by one or more R^{21} ;

R^{21} and R^{22} are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carboxy, carbamoyl, mercapto, sulphanoyl, trifluoromethyl, trifluoromethoxy,

methyl, ethyl, methoxy, ethoxy, vinyl, allyl, ethynyl, methoxycarbonyl, formyl, acetyl,

formamido, acetylamino, acetoxy, methylamino, dimethylamino, N -methylcarbamoyl,

N,N -dimethylcarbamoyl, methylthio, methylsulphonyl, mesyl, N -methylsulphanoyl and

N,N -dimethylsulphanoyl;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

10

2. A compound of formula (I) as claimed in claim 1 wherein one of R^1 and R^2 is ethyl and the other is butyl or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

15

3. A compound of formula (I) as claimed in either of claims 1 or 2 wherein R^3 is hydrogen or hydroxy or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

20

4. A compound of formula (I) as claimed in any one of claims 1-3 wherein v is 0 or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

5. A compound of formula (I) as claimed in any one of claims 1-4 wherein R^4 and R^6 are

hydrogen or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

25

6. A compound of formula (I) as claimed in any one of claims 1-5 wherein R^4 is hydrogen, halo or $C_{1-6}alkylS(O)_n$ wherein a is 0 or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

30

7. A compound of formula (I) as claimed in any one of claims 1-6 wherein R^5 is a group of formula (IA) (as depicted in claim 1); wherein

X is $-O-$;

Ring A is phenyl or 4-hydroxyphenyl;

R⁷ is hydrogen;

R⁸ is hydrogen;

R⁹ is hydrogen;

R¹¹ is carboxy; or R¹¹ is a group of formula (IB) (as depicted above); wherein:

R¹² is hydrogen;

R¹³ is hydrogen;

R¹⁵ is carboxy or sulpho;

p is 1 or 2;

q is 0;

r is 0;

m is 0; and

n is 1;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

8. A compound of formula (I) (as depicted above) wherein:

R¹ and R² are C₁₋₄alkyl;

v is 0;

R³ is hydrogen or hydroxy;

R³ and R⁶ are hydrogen;

R⁴ is hydrogen, halo or C₁₋₄alkylS(O)_n wherein a is 0;

R⁵ is a group of formula (IA) (as depicted above); wherein

X is -O-;

Ring A is aryl; wherein Ring A is optionally substituted by one or more substituents selected from R¹⁷;

R⁷ is hydrogen;

R⁸ is hydrogen;

R⁹ is hydrogen;

R¹¹ is carboxy; or R¹¹ is a group of formula (IB) (as depicted above); wherein:

R¹² is hydrogen;

R¹³ is hydrogen;

R¹⁵ is carboxy or sulpho;

p is 1 or 2;

q is 0;

r is 0;

m is 0;

n is 1; and

R¹⁷ is hydroxy;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

9. A compound of formula (I) selected from:

10 1,1-dioxo-3(R)-3-butyl-3-ethyl-5-(R)-5-phenyl-8-[N-(R)-α-carboxybenzyl]

carbamoylmethoxy]-2,3,4,5-tetrahydro-1,4-benzothiazepine;

1,1-dioxo-3(S)-3-butyl-3-ethyl-5-(S)-5-phenyl-8-[N-(R)-α-carboxybenzyl]

carbamoylmethoxy]-2,3,4,5-tetrahydro-1,4-benzothiazepine;

1,1-dioxo-3(R)-3-butyl-3-ethyl-5-(R)-5-phenyl-8-[N-(R)-α-[N-(carboxymethyl)carbamoyl]

benzyl]carbamoylmethoxy]-2,3,4,5-tetrahydro-1,4-benzothiazepine;

1,1-dioxo-3(S)-3-butyl-3-ethyl-5-(S)-5-phenyl-8-[N-(R)-α-[N-(carboxymethyl)carbamoyl]

benzyl]carbamoylmethoxy]-2,3,4,5-tetrahydro-1,4-benzothiazepine;

3,5-*trans*-1,1-dioxo-3-ethyl-3-butyl-5-phenyl-7-bromo-8-[N-(R)-α-[N-

(carboxymethyl)carbamoyl]benzyl]carbamoylmethoxy]-2,3,4,5-tetrahydro-1,4-

benzothiazepine;

3,5-*trans*-1,1-dioxo-3-(S)-3-ethyl-3-butyl-4-hydroxy-5-(S)-5-phenyl-7-bromo-8-[N-(R)-α-

[N-(carboxymethyl)carbamoyl]benzyl]carbamoylmethoxy]-2,3,4,5-tetrahydro-1,4-

benzothiazepine

3,5-*trans*-1,1-dioxo-3-(R)-3-ethyl-3-butyl-4-hydroxy-5-(R)-5-phenyl-7-bromo-8-[N-(R)-α-

[N-(carboxymethyl)carbamoyl]benzyl]carbamoylmethoxy]-2,3,4,5-tetrahydro-1,4-

benzothiazepine;

3,5-*trans*-1,1-dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-[N-(R)-α-[N-

(carboxymethyl)carbamoyl]benzyl]carbamoylmethoxy]-2,3,4,5-tetrahydro-1,4-

benzothiazepine;

3,5-*trans*-1,1-dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-[N-(R)-α-[N-(2-


sulphoethyl)carbamoyl]-4-hydroxybenzyl]carbamoylmethoxy]-2,3,4,5-tetrahydro-1,4-

benzothiazepine ammonium salt;

- 1,1-dioxo-3-(S)-3-ethyl-3-butyl-5-(S)-5-phenyl-7-methylthio-8-(N'-(R)- α -[N-(carboxymethyl)carbamoyl]benzyl)carbamoylmethoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine diethylamine salt; and
- 5 1,1-dioxo-3-(R)-3-ethyl-3-butyl-5-(R)-5-phenyl-7-methylthio-8-(N'-(R)- α -[N-(carboxymethyl)carbamoyl]benzyl)carbamoylmethoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine diethylamine salt;
- or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.


10. A process for preparing a compound of formula (I) or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as claimed in any one of claims 1-9, which process comprises of:

Process 1): oxidising a benzothiazepine of formula (II):




(II):

- 15 *Process 2*): for compounds of formula (I) wherein X is -O-, -NR^4 or -S-, reacting a compound of formula (IIIa) or (IIIb):



(IIIa)




(IIIb)


with a compound of formula (IV):

wherein L is a displaceable group;

Process 3): reacting an acid of formula (Va) or (Vb):




(Va)




(Vb)

or an activated derivative thereof, with an amine of formula (VI):




(VI):

- 10 *Process 4*): for compounds of formula (I) wherein R^11 is a group of formula (IB); reacting a compound of formula (I) wherein R^11 is carboxy with an amine of formula (VII):




(VII)

- 15 *Process 5*): for compounds of formula (I) wherein R^11 is a group of formula (IB) and R^15 is a group of formula (IC) reacting a compound of formula (I) wherein R^15 is carboxy with an amine of formula (VIII):



(VIII)

Process 6) for compounds of formula (I) wherein one of R^4 and R^5 are independently selected from C_{1-6} alkylthio optionally substituted on carbon by one or more R^{16} ; reacting a compound of formula (IXa) or (IXb):



(IXa)

(IXb)


wherein L is a displaceable group, with a thiol of formula (X):



(X)


wherein R^m is C_{1-6} alkylthio optionally substituted on carbon by one or more R^{16} ;

Process 7): for compounds of formula (I) wherein R^{11} is carboxy; deprotecting a compound of formula (XIa):




(XIa)

or (XIb):




(XIb)

wherein R^2 together with the $-OC(O)-$ group to which it is attached forms an ester; Process 8): for compounds of formula (I) wherein R^{11} is a group of formula (IB) and R^{15} is carboxy; deprotecting a compound of formula (XIIa):



(XIIa)

or (XIIb):




(XIIb)

- 62 -

wherein R^2 together with the $-OC(O)-$ group to which it is attached forms an ester,


Process 9): for compounds of formula (I) wherein R^{11} is a group of formula (IB) and Y is $-N(R^3)C(O)-$; reacting an acid of formula (XIIIa):



5


or (XIIIb):

(XIIIa)



(XIIIb)

or an activated derivative thereof, with an amine of formula (XIV):




(XIV)

or *Process 10):* for compounds of formula (I) wherein R^{11} is a group of formula (IB), R^{13} is a group of formula (IC) and R^{26} is carboxy, deprotecting a compound of formula (XVa):

- 63 -


or (XVb):

(XVa)



5

(XVb)



wherein R^2 together with the $-OC(O)-$ group to which it is attached forms an ester, and thereafter if necessary or desirable:

- i) converting a compound of the formula (I) into another compound of the formula (I);
- ii) removing any protecting groups;
- iii) forming a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug.

11. A compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as claimed in any one of claims 1 to 9 for use as a medicament.

12. A compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as claimed in any one of claims 1 to 9 for use in a method of prophylactic or therapeutic treatment of a warm-blooded animal, such as man.

13. The use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as claimed in any one of claims 1 to 9 in the manufacture of a medicament for use in the production of an IBAT inhibitory effect in a warm-blooded animal, such as man.

5

14. The use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as claimed in any one of claims 1 to 9, in the production of an IBAT inhibitory effect in a warm-blooded animal, such as man.

15. A method for producing an IBAT inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as claimed in any one of claims 1 to 9.

16. A pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as claimed in any one of claims 1 to 9, in association with a pharmaceutically-acceptable diluent or carrier.

17. A pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as claimed in any one of claims 1 to 9, and an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in association with a pharmaceutically acceptable diluent or carrier.

25

18. A pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as claimed in any one of claims 1 to 9, and a bile acid binder, in association with a pharmaceutically acceptable diluent or carrier.

30

19. A pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as

claimed in any one of claims 1 to 9, and an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a bile acid binder in association with a pharmaceutically acceptable diluent or carrier.

20. A composition according to claim 17 or claim 19 wherein the HMG Co-A reductase inhibitor is atorvastatin, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

21. A composition according to claim 17 or claim 19 wherein the HMG Co-A reductase inhibitor is rosuvastatin, or a pharmaceutically acceptable salt thereof.

22. A pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as claimed in any one of claims 1 to 9 and a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt thereof, in association with a pharmaceutically acceptable diluent or carrier.

23. A composition according to claim 22 wherein the PPAR alpha and/or gamma agonist is (S)-2-ethoxy-3-[4-(2-(4-methanesulphonyloxyphenyl)ethoxy)phenyl]propanoic acid or a pharmaceutically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/68 02/04043A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D281/10 A61K31/554 C07K5/06 A61K38/05 A61P3/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELD(S) SEARCHED
Mishpatim documentation searched (classification system followed by classification symbols)
IPC 7 C07D C07K

Documentation searched other than Mishpatim documentation to the extent that such documents are included in the lists searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Criterion of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 05188 A (MELLCOME FOUNDATION) 22 February 1996 (1996-02-22) cited in the application page 11, paragraph 6 -page 13, paragraph 5; claims; examples	1-23
A	WO 99 35135 A (ELAXO) 15 July 1999 (1999-07-15) claims; examples	1-23
P.X	WO 01 66533 A (ASTRAZENECA) 13 September 2001 (2001-09-13) page 1, line 1 -page 2, line 8; claims; examples	1-23

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

Special categories of cited documents:

- * document defining the general state of the art which is not considered to be of particular relevance
- * earlier document not published on or after the international filing date
- * document which may have priority claims or which may be cited in support of priority claims of another document relating to an oral disclosure, use, exhibition or other means
- * document published prior to the international filing date but later than the priority date claimed
- * document member of the same patent family

Data of the actual completion of the international search

Date of meeting of the international search report

11 November 2002

19/11/2002

Name and mailing address of the ISA
International Patent Office
IL - 2201 Tel Aviv, Israel
Tel: (+91-70) 540-2040, Tel. 51 651 6014
Fax: (+91-70) 540-2018

Authorized officer

Hejls, I

INTERNATIONAL SEARCH REPORT

International application No.
PCT/68 02/04043

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

The international Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:Because they relate to subject matter not required to be searched by the Authority, namely:
Although claim 15 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.2. ☐ Claims Nos.:

Because they relate to parts of the international Application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:

Because they are dependent claims and are not defined in accordance with the second and third sentences of Rule 8.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

The international Searching Authority found multiple inventions in the international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, the international Search Report covers all searchable claims.2. ☐ As all searchable claims could be searched without effort justifying an additional fee, the Authority did not invite payment of any additional fee.3. ☐ As only some of the required additional search fees were timely paid by the applicant, the international Search Report covers only those claims for which fees were paid, specifically claims Nos.:4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international Search Report is restricted to the invention first mentioned in the claims, it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

Form PCT/ISA210 (Continuation of first sheet (1)) (July 1989)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.
PCT/GB 02/04043

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9605188	A	22-02-1996	
AE	720 A	08-01-1999	
AU	696073 B2	03-09-1998	
AU	4426096 A	07-03-1996	
B6	62048 B1	29-01-1999	
B6	101209 A	29-08-1997	
BR	9508586 A	14-07-1998	
CA	2197099 A1	22-02-1996	
CN	1161035 A, B	01-10-1997	
CZ	9700373 A3	13-08-1997	
EE	9700028 A	15-08-1997	
EP	1203769 A1	08-05-2002	
EP	0775126 A1	28-05-1997	
FI	970531 A	07-02-1997	
MD	9605188 A1	22-02-1996	
HU	77129 A2	02-03-1998	
IL	114877 A	14-07-1999	
JP	2935756 B2	16-08-1999	
JP	10504035 T	14-04-1998	
NO	970585 A	07-04-1997	
NZ	290911 A	28-07-1998	
PL	318496 A1	23-06-1997	
RU	2156245 C2	20-09-2000	
SK	17797 A3	10-09-1997	
WO 9935135	A	15-07-1999	
AU	2515599 A	26-07-1999	
BR	9806799 A	10-10-2000	
CA	2317651 A1	15-07-1999	
CN	1292785 T	25-04-2001	
WO	9935135 A1	15-07-1999	
EP	1045840 A1	25-10-2000	
HR	20000468 A1	31-10-2000	
JP	2002500220 T	08-01-2002	
MD	20003514 A	07-09-2000	
PL	341672 A1	23-04-2001	
SK	1024200 A3	12-02-2001	
TR	200001816 T2	21-11-2000	
US	6465451 B1	15-10-2002	
ZA	9900081 A	06-07-2000	
WO 0166533	A	13-09-2001	
AU	3755601 A	17-09-2001	
WO	0166533 A1	13-09-2001	

BEST AVAILABLE COPY

THIS PAGE BLANK (USPTO)

THIS PAGE BLANK (USPTO)